

A survey of cryptosporidiosis among 2,541 residents of 25 coastal islands in Jeollanam-do (Province), Republic of Korea

Jae-Hwan PARK¹, Hyo-Jin KIM¹, Sang-Mee GUK¹, Eun-Hee SHIN¹, Jae-Lip KIM¹, Han-Jong RIM²,
Soon-Hyung LEE^{1,3} and Jong-Yil CHAI¹*

¹Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and
Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799,

²Department of Parasitology, College of Medicine, Korea University, Seoul, 136-705

³Korea Association of Health Promotion, Seoul 157-704, Korea

Abstract: In order to determine the distribution and prevalence of human cryptosporidiosis on western and southern coastal islands of Jeollanam-do (Province), fecal samples were collected from 2,541 people residing on 25 islands, 13 in the western coasts and 12 in the southern coasts, during July and August 2000. Fecal smears were prepared following formalin-ether sedimentation of the samples and stained by a modified acid-fast procedure. The presence of *Cryptosporidium* oocysts was determined by light microscopy. *Cryptosporidium* oocysts were detected in 38 specimens (1.5%). The oocyst positive rate varied (0-6.0%) according to island; the highest was detected on Oenarodo (6.0%), followed by Naenarodo (5.6%) and Nakwoldo (5.4%). The majority (35 persons, 94.6%) of *Cryptosporidium*-infected individuals were older than 50 years of age. Men (22/1,159; 1.9%) were infected at a higher rate than women (16/1,382; 1.2%). The results of the present survey indicate that human *Cryptosporidium* infections (due to *Cryptosporidium hominis* and/or *C. parvum*) are maintained at a relatively low prevalence on coastal islands of Jeollanam-do, Republic of Korea.

Key words: *Cryptosporidium hominis*, *Cryptosporidium parvum*, cryptosporidiosis, prevalence, coastal islands, Jeollanam-do (Province)

Cryptosporidium is a small coccidian protozoan which infects mammals, birds, reptiles, and fishes. The agents responsible for the majority of human infections are *Cryptosporidium hominis* (strong anthroponotic transmission) and *Cryptosporidium parvum* (zoonotic transmission) (Morgan-Ryan et al., 2002; Park et al., 2006).

In the Republic of Korea, after the first confirmation

of the presence of *C. parvum* in laboratory mice (Chai et al., 1990), high prevalences (according to detection methods: 10-22%) of human cryptosporidiosis were detected among out-patients at the Severance Hospital (Cho et al., 1993). Thereafter, cryptosporidiosis has been regarded as an important disease from the public health aspect in the Republic of Korea. Subsequently, several investigators have reported variable prevalences in differing localities (Chai et al., 1996; Seo et al., 2001; Yu et al., 2004; Park et al., 2006). In particular, epidemiological studies on human cryp-

• Received 12 October 2006, accepted after revision 8 November 2006.

*Corresponding author (e-mail: cji@snu.ac.kr)

Table 1. Prevalence of human cryptosporidiosis on 25 southern and western coastal islands of Jeollanam-do (Province), Republic of Korea

Location in Jeollanam-do	Name of island	Area code in Fig. 1	No. of positive / No. examined ^{a)} (%)
Southern coasts			
	Oenarodo	1	7 / 117 (6.0)
	Naenarodo	2	6 / 107 (5.6)
	Changdo	3	5 / 109 (4.6)
	Tolsando	4	1 / 93 (1.1)
	Nopdo	5	1 / 102 (1.0)
	Keogumdo	6	1 / 113 (0.9)
	Pyeongildo	7	0 / 108 (0.0)
	Kogumdo	8	0 / 111 (0.0)
	Sorokdo	9	0 / 91 (0.0)
	Pogildo	10	0 / 103 (0.0)
	Nohwado	11	0 / 109 (0.0)
	Sodo	12	0 / 100 (0.0)
Subtotal			21 / 1,263 (1.7)
Western coasts			
	Nakwoldo	13	5 / 93 (5.4)
	Hauido	14	4 / 96 (4.2)
	Cheungdo	15	3 / 92 (3.0)
	Hachodo	16	2 / 108 (1.9)
	Amtaedo	17	1 / 99 (1.0)
	Sangchodo	18	1 / 103 (1.0)
	Anchwado	19	1 / 110 (0.9)
	Kohado	20	0 / 85 (0.0)
	Dallido	21	0 / 104 (0.0)
	Tochodo	22	0 / 95 (0.0)
	Pikeumdo	23	0 / 104 (0.0)
	Imchado	24	0 / 99 (0.0)
	Heuksando	25	0 / 90 (0.0)
Subtotal			17 / 1,278 (1.3)
Total			38 / 2,541 (1.5)

^{a)}Examined by formalin-ether sedimentation and modified acid-fast (MAF) stain.

(4.2%), Cheungdo (3.0%), Hachodo (1.9%), Amtaedo (1.0%), Sangchodo (1.0%), and Anchwado (0.9%) (Table 1; Fig. 1). The average oocyst positive rate on the 13 western islands was 1.3% (17/1,278 examined). Among the total of 38 positive individuals, 22 (1.9%) were males and 16 (1.2%) were females (Fig. 2). The majority (94.6%) of the oocyst positive cases were over 50 years of age ($P = 0.000$) (Fig. 2).

In this epidemiological study, the overall human *Cryptosporidium* oocyst positive rate on the coastal

islands of Jeollanam-do, Korea was lower (1.5%) than expected. *Cryptosporidium* infections were detected on approximately half of the islands (13 of 25 islands). This relatively low prevalence of *Cryptosporidium* and the limited distribution on coastal islands of Jeollanam-do are interesting results. In previous studies conducted in inland areas of Jeollanam-do, higher prevalences were reported, 10.6% in 1992 (Chai et al., 1996) and 8.2% in 2002 (Yu et al., 2004). These discrepancies between coastal islands and inland areas




Fig. 2. The age- and sex-prevalence of cryptosporidiosis among people on the southern and western coastal islands of Jeollanam-do, Korea. Numbers (a/b) for each circle represent the number of oocyst positive cases (a) and the number of residents evaluated (b).

appear to be influenced by a variety of geographical and environmental factors.

Most coastal islands are geographically separated from inland areas. Migrations of people, as well as reservoir animals, between coastal islands and inland areas are less frequent than are observed within inland areas. Therefore, *Cryptosporidium* infection and transmission may be limited within islands.

Contamination of drinking water with oocysts constitutes an important risk factor for high cryptosporidiosis prevalence (Kelly et al., 1997). The islands examined in this study are geographically remote from inland areas, and the residents are unable to access treated water from inland. Thus, drinking water is supplied to the islanders from water reservoirs constructed within each island, or from underground. If water reservoirs were contaminated with *Cryptosporidium* oocysts, many populations would clearly be exposed to infection with the oocysts (Yamamoto et al., 2000). However, in the present study, we detected generally low cryptosporidiosis

prevalence on coastal islands, which may imply a relatively low possibility of water reservoir contamination. Another possibility includes the contamination of underground water. However, this possibility is also quite low, as the contamination of underground water with oocysts is uncommon and, on the majority of coastal islands, available water tends to contain some degree of salinity, which can negatively influence the viability of *Cryptosporidium* oocysts (El Mansoury et al., 2004).

In cases in which the probability of water contamination is low, contact with animal reservoirs should be considered. Domestic animals, including cattle and goats, represent important reservoir hosts for human cryptosporidiosis, and attendants for these animals are regularly exposed to the risk of cryptosporidiosis (Rahaman et al., 1984). However, it is regrettable that no studies have yet been published regarding the status of cryptosporidiosis in domestic animals on island areas in the Republic of Korea. In the case of inland Jeollanam-do, the *Cryptosporidium* oocyst positive rate among calves suffering from diarrhea was found to be 14.4% (Wee et al., 1996), and this rate was similar to that of humans (10.6%) occupying the same area (Chai et al., 1996). In another observation, the prevalence of cryptosporidiosis was surprisingly high (94%) among the domestic livestock reared in inland areas of Jeollanam-do (Yu et al., 2004).

Cryptosporidiosis induces mild, self-limiting diarrhea in healthy individuals, but can also cause severe diarrhea in young, elderly, or immunocompromised patients (Okhuysen et al., 1999). In the present study, it is worthy of note that the majority (94.6%) of oocyst positive cases were people over 50 years of age. In the questionnaire study, none of these people was determined to suffer from immunocompromised conditions or debilitating diseases. The majority of these patients complained only of mild degenerative diseases, which are common in subjects of advanced age. A similar predilection with regard to age has been reported in an inland rural village of Hwasun-gun, Jeollanam-do (Chai et al., 2001). This high prevalence among elderly subjects is distinct from other reports in other countries; human cryptosporidiosis appears

to predominate in children under 10 years of age (Esteban et al., 1998; Leach et al., 2000). One of the reasons for these differences includes that the majority of people living on the islands surveyed in the present study were adults of over 30 years of age and very few children were living on these islands.

Seasonal variations of human cryptosporidiosis have also been reported (McLauchlin et al., 2000; Chai et al., 2001). In the Republic of Korea, as the result of monthly observations in an inland village of Jeollanam-do, the *Cryptosporidium* oocyst positive rate was lower during summer than in the spring and autumn seasons (Chai et al., 2001). The present study was conducted during the summer season, from July to August. In order to elucidate the seasonal tendencies on the islands surveyed herein, further studies will be necessary.

Unfortunately, we could not isolate *Cryptosporidium* DNA, due to the insufficient amount of fecal samples and of oocysts collected from each of the individuals assessed in this study. Therefore, we were unable to analyze the genotype and species of *Cryptosporidium* present on these islands. Our speculation, although unsupported by evidence, is that the infections may represent a mixed population of *C. hominis* and *C. parvum*, as has been previously observed in an inland area of Jeollanam-do (Park et al., 2006).

In conclusion, the status of human cryptosporidiosis on the coastal islands of Jeollanam-do, Republic of Korea, has been studied in this work for the first time. It has been determined that these islands evidence a relatively low and variable prevalence (0-6.0%) of cryptosporidiosis. The species of *Cryptosporidium* involved, i.e., *C. hominis*, *C. parvum*, or both, transmission modes, and other epidemiological characteristics of cryptosporidiosis on these islands remain to be studied in the future.

ACKNOWLEDGMENTS

The authors are indebted to members of the headquarters and branch offices of the Korea Association of Health Promotion for their assistance in the collection of the fecal specimens.

REFERENCES

- Chai JY, Shin SM, Yun CK, Yu JR, Lee SH (1990) Experimental activation of cryptosporidiosis in mice by immunosuppression. *Korean J Parasitol* **28**: 31-37.
- Chai JY, Lee SH, Guk SM, Lee SH (1996) An epidemiological survey of *Cryptosporidium parvum* infection in randomly selected inhabitants of Seoul and Chollanam-do. *Korean J Parasitol* **34**: 113-119.
- Chai JY, Kim NY, Guk SM, Park YK, Seo M, Han ET, Lee SH (2001) High prevalence and seasonality of cryptosporidiosis in a small rural village occupied predominantly by aged people in the Republic of Korea. *Am J Trop Med Hyg* **65**: 518-522.
- Chai JY, Lee SH (2002) Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* **51**: 129-154.
- Cho MH, Kim AK, Im KI (1993) Detection of *Cryptosporidium* oocysts from out-patients of the Severance Hospital, Korea. *Korean J Parasitol* **31**: 193-199.
- El Mansoury ST, Abou El Naga IF, Negm AY, Amer EE (2004) Influence of temperature and salinity on the viability and infectivity of *Giardia lamblia* and *Cryptosporidium parvum*. *J Egypt Soc Parasitol* **34**: 161-172.
- Esteban JG, Aguirre C, Flores A, Strauss W, Angles R, Mas-Coma S (1998) High *Cryptosporidium* prevalences in healthy Aymara children from the northern Bolivian Altiplano. *Am J Trop Med Hyg* **58**: 50-55.
- Goo GS, Min DY, Ahn MH, Kim KM, Leem MH, Yoon HS (1988) Status of intestinal parasitic infections in a remote island, Yondo, Jeonranam-do: province. *Korean J Parasitol* **26**: 275-284.
- Kelly P, Baboo KS, Ndubani P, Nchito M, Okeowo NP, Luo NP, Feldman RA, Farthing MJ (1997) Cryptosporidiosis in adults in Lusaka, Zambia, and its relationship to oocyst contamination of drinking water. *J Infect Dis* **176**: 1120-1123.
- Leach CT, Koo FC, Kuhls TL, Hilsenbeck SG, Jenson HB (2000) Prevalence of *Cryptosporidium parvum* infection in children along the Texas-Mexico border and associated risk factors. *Am J Trop Med Hyg* **62**: 656-661.
- McLauchlin J, Amar C, Pedraza-Diaz S, Nichols GL (2000) Molecular epidemiological analysis of *Cryptosporidium* spp. in the United Kingdom: results of genotyping *Cryptosporidium* spp. in 1,705 fecal samples from humans and 105 fecal samples from livestock animals. *J Clin Microbiol* **38**: 3984-3990.
- Morgan-Ryan UM, Fall A, Ward LA, Hijjawi N, Sulaiman I, Fayer R, Thompson RC, Olson M, Lal A, Xiao L (2002)

- Cryptosporidium hominis* n. sp. (Apicomplexa: Cryptosporidiidae) from *Homo sapiens*. *J Eukaryot Microbiol* **49**: 433-440.
- Okhuysen PC, Chappell CL, Crabb JH, Sterling CR, DuPont HL (1999) Virulence of three distinct *Cryptosporidium parvum* isolates for healthy adults. *J Infect Dis* **180**: 1275-1281.
- Park JH, Guk SM, Han ET, Shin EH, Kim JL, Chai JY (2006) Genotype analysis of *Cryptosporidium* spp. prevalent in a rural village in Hwasun-gun, Republic of Korea. *Korean J Parasitol* **44**: 27-33.
- Rahaman AS, Sanyal SC, Al-Mahmud KA, Sobhan A, Hossain KS, Anderson BC (1984) Cryptosporidiosis in calves and their handlers in Bangladesh. *Lancet* **28**: 221.
- Seo M, Huh S, Chai JY, Yu JR (2001) An epidemiology survey on *Cryptosporidium parvum* infection of inhabitants in Chorwon-gun, Kangwon-do. *Korean J Parasitol* **39**: 201-203.
- Wee SH, Joo HD, Kang YB (1996) Evaluation for detection of *Cryptosporidium* oocysts in diarrheal feces of calves. *Korean J Parasitol* **34**: 121-126.
- Yamamoto N, Urabe K, Takaoka M, Nakazawa K, Gotoh A, Haga M, Fuchigami H, Kimata I, Iseki M (2000) Outbreak of cryptosporidiosis after contamination of the public water supply in Saitama Prefecture, Japan, in 1996. *Kansenshogaku Zasshi* **74**: 518-526.
- Yu JR, Lee JK, Seo M, Kim SI, Sohn WM, Huh S, Choi HY, Kim TS (2004) Prevalence of cryptosporidiosis among the villagers and domestic animals in several rural areas of Korea. *Korean J Parasitol* **42**: 1-6.