

Foodborne Intestinal Flukes in Southeast Asia

Jong-Yil Chai^{1,*}, Eun-Hee Shin¹, Soon-Hyung Lee¹ and Han-Jong Rim²

¹Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799, Korea; ²Department of Parasitology, College of Medicine, Korea University, Seoul 136-705, Korea

Abstract: In Southeast Asia, a total of 59 species of foodborne intestinal flukes have been known to occur in humans. The largest group is the family Heterophyidae, which constitutes 22 species belonging to 9 genera (*Centrocestus*, *Haplorchis*, *Heterophyes*, *Heterophyopsis*, *Metagonimus*, *Procerovum*, *Pygidiopsis*, *Stellantchasmus*, and *Stictodora*). The next is the family Echinostomatidae, which includes 20 species in 8 genera (*Artyfechinostomum*, *Acanthoparyphium*, *Echinochasmus*, *Echinoparyphium*, *Echinostoma*, *Episthmium*, *Euparyphium*, and *Hypoderaeum*). The family Plagiorchiidae follows the next containing 5 species in 1 genus (*Plagiorchis*). The family Lecithodendriidae includes 3 species in 2 genera (*Phaneropsolus* and *Prosthodendrium*). In 9 other families, 1 species in 1 genus each is involved; Cathaemaciidae (*Cathaemacia*), Fasciolidae (*Fasciolopsis*), Gastrodiscidae (*Gastrodiscoides*), Gymnophallidae (*Gymnophalloides*), Microphallidae (*Spelotrema*), Neodiplostomidae (*Neodiplostomum*), Paramphistomatidae (*Fischoederius*), Psilostomidae (*Psilorchis*), and Strigeidae (*Cotylurus*). Various types of foods are sources of human infections. They include freshwater fish, brackish water fish, fresh water snails, brackish water snails (including the oyster), amphibians, terrestrial snakes, aquatic insects, and aquatic plants. The reservoir hosts include various species of mammals or birds. The host-parasite relationships have been studied in *Metagonimus yokogawai*, *Echinostoma hortense*, *Fasciolopsis buski*, *Neodiplostomum seoulense*, and *Gymnophalloides seoi*; however, the pathogenicity of each parasite species and host mucosal defense mechanisms are yet poorly understood. Clinical aspects of each parasite infection need more clarification. Differential diagnosis by fecal examination is difficult because of morphological similarity of eggs. Praziquantel is effective for most intestinal fluke infections. Continued efforts to understand epidemiological significance of intestinal fluke infections, with detection of further human cases, are required.

Key words: intestinal fluke, foodborne, heterophyid, echinostome, epidemiology, Asia

INTRODUCTION

Although 40-50 million people are generally estimated to be infected with foodborne intestinal trematodes worldwide [1], this certainly is an underestimate of the true number of people infected. Most of the infected people live in Southeast Asia, including Korea, China, Thailand, Vietnam, Lao PDR, the Philippines, Indonesia, and India. The number of trematode species, currently known to be involved in this area, is counted at least 59, including *Metagonimus yokogawai*, *Heterophyes nocens*, *Haplorchis taichui*, *Prosthodendrium molenkampi*, *Phaneropsolus bonnei*, *Echinostoma hortense*, *Echinochasmus japonicus*, *Fasciolopsis buski*, *Neodiplostomum seoulense*, and *Gymnophalloides seoi* [1-6] (Table 1). They belong to the families Heterophyidae, Echinostomatidae Poche, 1926, Plagiorchiidae Ward, 1917, Lecithodendriidae

Odhner, 1911, Neodiplostomidae Shoop 1989, Paramphistomatidae Fischoeder, 1901, Cathaemaciidae Fuhrmann, 1928, Fasciolidae Railliet, 1895, Gastrodiscidae Stiles and Goldberger, 1910, Gymnophallidae Morozov, 1955, Microphallidae Trava-sos, 1920, Psilostomidae Odhner, 1913, and Strigeidae Railliet, 1919 [1-6]. In this paper, morphology, biology, epidemiology, pathogenicity, clinical manifestations, diagnosis, and treatment of these fluke infections are briefly reviewed.

SPECIES INFECTING HUMANS

Acanthoparyphium tyosenense Yamaguti, 1939

This echinostome is morphologically characterized by the presence of a characteristic head collar with a total of 23 collar spines arranged in a single row, without ventral corner spines, a long cirrus sac reaching beyond the posterior margin of the acetabulum, and the vitellaria extending to the level of the cirrus sac or the Mehlis' gland [7]. This species was originally described based on worms from the small intestines of the duck

• Received 28 September 2009, revised 5 October 2009, accepted 5 October 2009.

* Corresponding author (cyj@snu.ac.kr)

Table 1. Foodborne intestinal flukes and the infection source in Southeast Asia

Parasite species	Source of human or animal infections
<i>Acanthoparyphium tyosenense</i>	Bivalves, <i>Macra veneriformis</i> , <i>Solen grandis</i> ; gastropod, <i>Neverita bicolor</i>
<i>Artyfechinostomum malayanum</i>	Snail, <i>Digoniostoma pulchella</i> <i>Echinostoma malayanum</i> , <i>Pila scutata</i> , <i>Lymnaea (Bullastra) cumingiana</i>
<i>Artyfechinostomum oraoni</i>	Probably freshwater snail
<i>Ascocotyle (Phagicola) longa</i>	Freshwater fish
<i>Cathaemacia cabrerai</i>	Unknown
<i>Centrocestus armatus</i>	Fresh water fish, <i>Zacco platypus</i> , <i>Zacco temminckii</i> , <i>Rhodeus ocellatus</i> , <i>Gobius similis</i> , <i>Pseudorasbora parva</i> , <i>Pelteobagrus fulvidraco</i>
<i>Centrocestus caninus</i>	Freshwater fish, <i>Cyprinus carpio</i> , <i>Hampala dispar</i> , <i>Puntius</i> spp., <i>Cyclocheilichthys</i> sp., <i>Tilapia nilotica</i>
<i>Centrocestus formosanus</i>	Freshwater fish
<i>Centrocestus kurokawai</i>	Probably freshwater fish
<i>Cotylurus japonicus</i>	Freshwater snail, <i>Stagnicola</i> , <i>Lymnaea</i> , <i>Physa</i> , <i>Heligsoma</i> spp.
<i>Cryptocotyle lingua</i>	Freshwater fish, <i>Gobius ruthensparri</i> , <i>Labrus bergylta</i>
<i>Echinochasmus fujianensis</i>	Freshwater fish, <i>Pseudorasbora parva</i> , <i>Cyprinus carpio</i>
<i>Echinochasmus japonicus</i>	Fresh water fish, <i>Pseudorasbora parva</i> , <i>Hypomesus olidus</i> , <i>Gnathopogon strigatus</i>
<i>Echinochasmus jiufoensis</i>	Probably freshwater fish
<i>Echinochasmus liliputanus</i>	Freshwater fish, <i>Pseudorasbora parva</i> , goldfish
<i>Echinochasmus perfoliatus</i>	Freshwater fish, <i>Carassius</i> sp.
<i>Echinoparyphium recurvatum</i>	Freshwater snail, <i>Planorbis planorbis</i> , <i>Lymnaea</i> sp., <i>Lymnaea stagnalis</i> ; tadpole and frog of <i>Rana temporaria</i>
<i>Echinostoma angustitestis</i>	Freshwater fish
<i>Echinostoma cinetorchis</i>	Freshwater fish, <i>Misgurnus anguillicaudatus</i> ; freshwater snail, <i>Radix auricularia coreanus</i> , <i>Physa acuta</i> , <i>Cipangopaludina chinensis malleata</i>
<i>Echinostoma echinatum</i>	Mussel, <i>Corbicula lindoensis</i> , <i>Corbicula sucplanta</i> , <i>Idiopoma javanica</i> ; freshwater snail, <i>Biomphalaria glabrata</i>
<i>Echinostoma hortense</i>	Freshwater fish, <i>Misgurnus anguillicaudatus</i> , <i>Misgurnus mizolepis</i> , <i>Odontobutis obscura interrupta</i> , <i>Moroco oxycephalus</i> , <i>Coreoperca kawamebari</i> , <i>Squalidus coreanus</i>
<i>Echinostoma ilocanum</i>	Large snail, <i>Pila conica</i> , <i>Viviparus javanicus</i>
<i>Echinostoma macrorchis</i>	Large snail, <i>Cipangopaludina malleata</i> , <i>Cipangopaludina japonica</i> , <i>Segmentina nitiella</i> , <i>Viviparus malleatus</i> ; frog of <i>Rana</i> sp.
<i>Echinostoma revolutum</i>	Snail or clam, <i>Corbicula product</i> ; tadpole
<i>Episthmium caninum</i>	Freshwater fish
<i>Fasciolopsis buski</i>	Aquatic plant, including water caltrop, water cress, water chestnut, and water bamboo
<i>Fischoederius elongatus</i>	Aquatic plant
<i>Gastrodiscoides hominis</i>	Tadpole, frog, crayfish, aquatic plant
<i>Gymnophalloides seoi</i>	Oyster, <i>Crassostrea gigas</i>
<i>Haplorchis pumilio</i>	Freshwater fish, Cyprinidae, Siluridae, Cobitidae
<i>Haplorchis taichui</i>	Freshwater fish, <i>Cyprinus carpio</i> , <i>Carassius auratus</i> , <i>Zacco platypus</i> , <i>Pseudorasbora parva</i> , <i>Rodeus ocellatus</i> , <i>Gambusia affinis</i> , <i>Puntius orphoides</i> , <i>Puntius</i> spp., <i>Raiamas guttatus</i> , <i>Mystacoleucus marginatus</i> , <i>Henichoryhnchus siamensis</i>
<i>Haplorchis vanissimus</i>	Freshwater fish
<i>Haplorchis yokogawai</i>	Freshwater fish, <i>Mugil</i> spp., <i>Puntius</i> spp., <i>Misgurnus</i> sp., <i>Ophicephalus striatus</i>
<i>Heterophyes nocens</i>	Brackish water fish, <i>Mugil</i> sp., <i>Acanthogobius</i> sp.
<i>Heterophyopsis continua</i>	Brackish water fish, <i>Acanthogobius</i> sp., <i>Lateolabrax</i> sp., <i>Clupadon punctatus</i>
<i>Hypoderaeum conoideum</i>	Snail, tadpole
<i>Isthmiophora melis</i>	Tadpole, loach <i>Misgurnus</i> spp.
<i>Metagonimus minutus</i>	Mullet, <i>Mugil cephalus</i>
<i>Metagonimus miyatai</i>	Sweetfish, dace, fat-minnow <i>Morocco steindachneri</i> , pale chub <i>Zacco platypus</i> , dark chub <i>Zacco temmincki</i> , Crussian carp <i>Carrassius carassius</i> , carp <i>Cyprinus carpio</i> , dace <i>Tribolodon taczanowskii</i> , perch <i>Lateolabrax japonicus</i>
<i>Metagonimus takahashii</i>	
<i>Metagonimus yokogawai</i>	Sweetfish <i>Plecoglossus altivelis</i> , dace <i>Tribolodon</i> sp., perch <i>Lateolabrax japonicus</i>
<i>Neodiplostomum seoulense</i>	Grass snake, <i>Rhabdophis tigrina</i> ; tadpole and frog of <i>Rana nigromaculata</i>
<i>Phaneropsolus bonnei</i>	Naiad of dragonfly, damselfly
<i>Phaneropsolus spinicirrus</i>	Probably naiads of dragonfly
<i>Plagiorchis harinasutai</i>	Probably insect larva
<i>Plagiorchis javensis</i>	Insect larva
<i>Plagiorchis muris</i>	Insect larval, insect naiad
<i>Plagiorchis philippinensis</i>	Insect larva

(Continued to the next page)

Table 1. (Continued from the previous page)

Parasite species	Source of human or animal infections
<i>Plagiorchis muris</i>	Freshwater fish, various species, freshwater snail
<i>Plagiorchis vespertilionis</i>	Mosquito larva, caddis-fly larva, mayfly larva, dragonfly nymph; freshwater fish
<i>Procerovum calderoni</i>	Freshwater fish, <i>Ophiocephalus striatus</i> , <i>Glossogobius giurus</i> , <i>Mollienesia latipinna</i> , <i>Mugil</i> sp., <i>Creisson validus</i>
<i>Procerodum varium</i>	Mullet, <i>Mugil cephalus</i>
<i>Prosthodendrium molencampi</i>	Naiad of dragonfly, damselfly
<i>Psilorchis hominis</i>	Unknown
<i>Pygidiopsis summa</i>	Mullet <i>Mugil cephalus</i> , goby <i>Acanthogobius flavimanus</i>
<i>Spelotrema brevicacaeca</i>	Crab <i>Carcinus maenas</i> ; shrimp <i>Macrobrachium</i> sp.
<i>Stellantchasmus falcatus</i>	Mullet and half-beaked fish <i>Dermogenus pusillus</i>
<i>Stellantchasmus formosanus</i>	Mullet <i>Mugil cephalus</i>
<i>Stellantchasmus pseudocirratus</i>	Mullet <i>Mugil cephalus</i>
<i>Stictodora fuscata</i>	Goby <i>Acanthogobius flavimanus</i>
<i>Stictodora lari</i>	Goby <i>Acanthogobius flavimanus</i> , other estuarine fish

Melanitta fusca stejnegeri and *Melanitta nigra americana* caught in the Republic of Korea [8]. This echinostome is at present known to be distributed in the Republic of Korea and Japan [9]. The first intermediate hosts include marine megagastropods *Lunatia fortuni* and *Glossaulax didyma* in the Republic of Korea [9]. *Cercaria yamagutii* shed from marine gastropods, *Tympanotonus microptera*, *Cerithidea cingulata*, and *Cerithidea largillierti*, in Japan [10], is now considered to be the cercariae of *A. tyosenense* [9]. Four species of brackish water bivalves, i.e., *Macra veneriformis*, *Solen grandis*, *Solen strictus*, and *Ruditapes philippinarum*, and a species of brackish water gastropod, *Neverita bicolor*, have been verified to be the second intermediate hosts [7,9]. Chicks [7,11] and sea gulls *Larus crassirostris* [9] were reported to be the experimental definitive hosts. In chicks, worms grew quickly to become ovigerous adults in the jejunum within 5 days and survived at least up to 38 days [11]. The worms fully grew to have a peak number of uterine eggs during day 10 and day 15 after infection [11]. Ten human infections were first discovered in 2 coastal villages of Chollabuk-do Province, Republic of Korea [7]. The patients used to eat improperly cooked marine bivalves and gastropods.

Artyfechinostomum malayanum (Leiper, 1911) Mendheim, 1943

(syn. *Echinostoma malayanum* Leiper, 1911; *Euparyphium malayanum* Odhner, 1913; *Artyfechinostomum sufrartifex* Lane, 1915; *Artyfechinostomum mehrai* Faruqui, 1930; *Paryphostomum sufrartifex* Bhalerao, 1931; *Isthmiophora malayana* [48])

This echinostome was first described from a human in Malaysia under the name *Echinostoma malayanum* Leiper, 1911 [12], and then in Singapore, Thailand, Indonesia, India, and the Philippines [6,13-15]. It has a small head collar, with a total of 43 (39-45) collar spines arranged in 2 alternating rows at the dor-

sal side, 2 deeply lobed testes, and a large and long cirrus sac reaching beyond the ventral sucker [16]. It was noted that *E. malayanum* Leiper, 1911 fits better to the generic features of *Artyfechinostomum* Lane, 1915 [16,17]. A closely related species, *A. sufrartifex*, was found in an Assamese girl [18] and a boy [19] in India, and also in pigs [20], and cats and dogs [21]. *A. sufrartifex* (Lane, 1915) was found conspecific with *A. malayanum* (Leiper, 1911) [17]. Another species, *A. mehrai* Faruqui, 1930, which was later synonymized with *A. malayanum* [17,20], was described in 1930 from a Hindu girl suffering from diarrhea, vomiting, anorexia, and loss of weight [22]. *A. mayananum* is now known to be distributed in Malaysia, Thailand, Indonesia, India, and the Philippines [6,12-15]. The first intermediate host is the freshwater snail, *Indoplanorbis exustus* or *Gyraulus convexiusculus*, and cercariae encyst in various species of snails, i.e., *Pila scutata*, *Lymnaea (Bullastra) cumingiana*, and *Digoniostoma pulchella* [6]. Humans, pigs, rats, cats, dogs, mice, hamsters, and house-shrews were reported to be the definitive hosts [6,16,21,23].

Artyfechinostomum oraoni Bandyopadhyay, Manna & Nandy, 1989

This echinostome was first reported from 20 human infections in a tribal community near Calcutta, India [24,25]. The first intermediate host is the freshwater snail, *Lymnaea* sp. [26]. In naturally infected pigs, *A. oraoni* provoked fatal diarrhea [25].

Cathaemacia cabrerai Jueco & Monzon, 1984

This fluke was first reported from a patient in the Philippines [28]. No further information is available.

Centrocestus armatus (Tanabe, 1922) Price, 1932

This heterophyid fluke, having 42-48 circumoral spines, was

described from experimental dogs, cats, rabbits, rats, and mice fed on cyprinoid fish infected with the metacercariae [29]. Characteristic morphological features include a small number of intrauterine eggs, the median location of the ovary, and the side-by-side location of the 2 testes [30]. An experimental human infection was reported successful in Japan [29], and a case of natural human infection was reported in the Republic of Korea [30]. The first intermediate host is the fresh water snail, *Semisulcospira* sp. [31], and the second intermediate hosts are freshwater fish, including *Zacco platypus*, *Zacco temminckii*, *Rhodeus ocelatus*, *Gobius similis*, *Pseudorasbora parva*, and *Pelteobagrus fulvidraco* [32,33]. The natural definitive hosts include the large egret *Egretta alba modesta* [34] and the cat [35].

Centrocestus caninus (Leiper, 1913) Yamaguti, 1958

(syn. *Stephanopirum longus* Onji & Nishio, 1916)

This heterophyid fluke, armed with 26-30 circumoral spines, was originally described in 1913 based on a single specimen obtained from a dog in Taiwan [4,36], and 2 human infections were diagnosed with this species in Thailand [37]. The freshwater snail, *Melanoides tuberculata*, is the snail host, and freshwater fish, such as, *Carrasius auratus*, *Cyprinus carpio*, *Hampala dispar*, *Puntius* sp., *Cyclocheilichthys* sp., and *Tilapia nilotica*, serve as the second intermediate hosts [37,38]. The validity of this species is now questioned, and this species seems synonymous with *C. formosanus* [36].

Centrocestus formosanus (Nishigori, 1924) Price, 1932

This heterophyid fluke, having 32 circumoral spines, was originally described from an experimental dog fed freshwater fish infected with the metacercariae [39]. An experimental human infection and a natural infection in a fox were also reported [39]. Natural human infections were found in Taiwan and Japan [40, 41]. The distribution of this fluke is known in Taiwan, China, Japan, Philippines, India, Hawaii, Mexico, and Lao PDR [6,36, 41-45]. The snail, *Stenomelania newcombi*, sheds the cercariae [43]. Freshwater fish, including *Cyclocheilichthys repasson*, *Puntius brevis*, and *Osteochilus hasseltii*, harbor the metacercariae [36,46].

Centrocestus kurokawai (Kurokawa, 1935) Yamaguti, 1958

This heterophyid fluke, having 38-40 circumoral spines, was described from a naturally infected man in Hiroshima Prefecture, Japan [47]. No other information is available on its life cycle.

Cotylurus japonicus Ishii, 1932

This strigeid fluke was originally described from ducks and chickens in Japan by Ishii in 1932 [48]. Later it was reported from a 13-year-old girl in Hunan Province, China [49]. Freshwater snails belonging to the genera *Stagnicola*, *Lymnaea*, *Physa*, and *Heligsoma* shed the cercariae, and the cercariae encyst in the same snail hosts to become specialized metacercariae known as 'tetracotyles' [1]. Human infections may occur when tetracotyles in infected snails are ingested [1].

Echinochasmus fujianensis Cheng et al., 1992

This echinostome was originally described from man, dogs, cats, pigs, and rats in Fujian Province, China [50]. The head collar is prominent with 24 spines arranged in a single row, interrupted dorsally [50]. Genetically *E. fujianensis* is distinguished from *Echinochasmus japonicus* by means of random amplified polymorphic DNA analysis (RAPD) patterns [51]. In 5 areas of southern Fujian Province, the prevalence among residents was 3.2% (1.6-7.8% in range), and about two-thirds of the infected people were 3-15 years of age [6,50]. Among the 3 *Echinochasmus* species, *E. fujianensis*, *E. japonicus*, and *E. perfoliatus*, existing in Hubei, Anhui, and Fujian provinces, China, *E. fujianensis* is the dominant species [51]. Freshwater snails, *Bellamya aeruginosa*, shed the cercariae, and *Pseudorasbora parva* and *Cyprinus carpio* harbor the metacercariae [6]. Dogs, cats, pigs, and rats serve as the natural definitive hosts [50].

Echinochasmus japonicus Tanabe, 1926

This echinostome, having dorsally interrupted 24 collar spines, was first described in Japan from experimental animals, including dogs, cats, rats, mice, and birds fed the metacercariae encysted in freshwater fish [52], and is now known to occur in the far East [2,5]. Its morphological characters include a small and plump body, 2 large and tandem testes, and a very small number of eggs, usually less than 5, in the uterus [2]. An successful experimental human infection was reported in Japan [53], and later natural human infections were found in several provinces of China [54,55] and the Republic of Korea [56]. Fresh water snails, *Parafossarulus manchouricus*, shed the cercariae [57,58], and 18 species of freshwater fish, including *Pseudorasbora parva*, *Hypomesus olidus*, and *Gnathopogon strigatus*, harbor the metacercariae [58-60]. In the laboratory, its life cycle from cercariae to adults has been successfully completed [58]. Avian species, such as ducks [61] and egrets [34], and mammalian species, such as cats [35] and insectivores [62], have been confirmed as

the natural definitive hosts. This echinostome is prevalent among people, as well as animals, in 6 counties of Fujian and Guangdong, China, the prevalence being 4.9% in humans, 39.7% in dogs, and 9.5% in cats [6].

Echinochasmus jiufuensis Yu & Mott, 1994

This echinostome, having dorsally interrupted 24 collar spines, was first described in 1988 at an autopsy of a 6-month-old girl who died from pneumonia and dehydration in Guangzhou, China [63]. This fluke closely resembles *Echinochasmus beleocephalus* Dietz, 1909, but distinguished in several points [6,63]. The life cycle is unknown [63].

Echinochasmus liliputanus (Looss, 1896) Odhner, 1910

This echinostome, having dorsally interrupted 24 collar spines, was originally reported from dogs, cats, and birds in Egypt, Syria, and Palestine [48]. The head collar is reniform, and the vitellarium does not extend anterior to the ventral sucker [64]. Human infections were first discovered in Anhui Province, China in 1991, with the prevalence rate of 13.4% among 2,426 people [65]. The infection rates in dogs and cats living in the same place were 60% and 45%, respectively [65]. Thereafter, more than 2,500 human infection cases have been reported in Anhui Province, China [66]. The freshwater snail *Parafossarulus striatulus* shed the cercariae [6], and the freshwater fish *Pseudorasbora parva* [6] and goldfish [66] harbor the metacercariae. Badgers, foxes, raccoons, dogs, and cats serve as the natural definitive hosts [6,67]. Humans may be infected with this parasite through drinking untreated water containing the cercariae [68]. Cercariae were found to be encysted in the presence of human gastric juice, and this was proposed as the plausible mechanism of cercariae-induced infections [66].

Echinochasmus perfoliatus (Ratz, 1908) Gedoelst, 1911

This echinostome, having dorsally interrupted 24 collar spines, was first recovered from dogs in Rumania by Motas and Straulescus in 1902 under the name *Distoma echinatum*, and then found again by Ratz in 1908 from dogs and cats in Hungary [69]. It is now a common intestinal parasite of dogs and cats in Hungary, Italy, Rumania, Russia, Japan, China, and Taiwan [6, 70], and of red foxes in Denmark [71]. Its body is elongated and larger than the closely related species, *E. japonicus*. An experimental [69] and a natural human infection [72] were successively reported in Japan. A low prevalence (1.8%) of human infections was reported from Guangdong, Fujian, Anhui, and

Hubei Provinces of China; however, a child died from the infection from whom about 14,000 worms were recovered at autopsy [6]. Freshwater snails, *Parafossarulus manchouricus*, *Bithynia leachi*, and *Lymnaea stagnalis* shed the cercariae [48]. Freshwater fish, such as *Carassius* sp., *Zacco platypus*, *Zacco teminckii*, and *Pseudorasbora parva*, harbor the metacercariae, which are encysted only on the gills [6,73]. Mammals, including rats, cats, dogs, foxes, fowls, and wild boars are the reservoir hosts [12,71,74].

Echinoparyphium recurvatum (von Linstow, 1873) Lühe, 1909

(syn. *Echinoparyphium koidzumii* Tsuchimochi, 1924)

This echinostome was first found in various avian species, and now also recognized as a parasite of mammals [6,12,75,76]. The first human infection was identified in Taiwan in 1929 and in Indonesia in 1948 [12]. It is characterized by having a relatively small body, a head collar armed with 45 collar spines arranged in 2 alternating rows and 4-5 end group spines on each side, a large ventral sucker, and a short uterus with only a few eggs [73,75]. It has been suggested that *E. recurvatum* complex consists of at least 3 species, including 2 other similar species, i.e., *E. pseudorecurvatum* and *E. mordwilkoii* [77]. The planorbisid snails, including *Physa alexandrina*, *Physa fontinalis*, *Planorbis planorbis*, *Lymnaea pervia*, *Lymnaea peregra*, *Valvata piscinalis*, and *Radix auricularia coreana* shed the cercariae [78-80], and the metacercariae encyst in tadpoles and frogs of *Rana temporaria* and also in snails, *P. planorbis*, *Lymnaea* sp. [6], *R. auricularia coreana* [80], and *Lymnaea stagnalis* [81]. House rats [75], wild rats (*Arvicanthis niloticus*) [82], and species of birds are natural definitive hosts [48]. Human infections have been recorded in Taiwan, Indonesia, and Egypt [6,12].

Echinostoma angustitestis Wang, 1977

This echinostome was first described from dogs experimentally infected with metacercariae isolated from the freshwater fish in 1977 in China [6]. It has a head collar armed with a total of 41 collar spines arranged in 2 alternating rows [83]. Only 2 human infections were reported in Fujian, China, who complained of dizziness, abdominal pain, and diarrhea [83]. There is no available information on its life history.

Echinostoma cinetorchis Ando & Ozaki, 1923

This echinostome was first discovered in rats in Japan [84], and then in a dog in Taiwan [85,86], rats in the Republic of Korea [87,88]. This fluke also exists in China [89]. Human infection

cases were first reported in Japan [90,91] and then in the Republic of Korea [92-94]. Its head collar is equipped with total 37-38 spines and abnormal location and/or disappearance of one or both testes is a characteristic feature [2]. The freshwater snails *Hippeutis cantori* [95] and *Segmentina hemisphaerula* [96] shed the cercariae and also harbor the metacercariae [95]. Other freshwater snails, including *Radix auricularia coreana*, *Physa acuta*, *Cipangopaludina* sp., and *Cipangopaludina chinensis malleata*, were proved to be the second intermediate hosts [97,98]. Tadpoles of *Rana nigromaculata*, *Rana rugosa*, and *Rana japonica* [99], and freshwater fish, especially the loach *Misgurnus anguillicaudatus* [100,101], were proven to harbor the metacercariae. Rats [62,84, 87,88] and dogs [102] serve as the natural definitive hosts. Rats and mice are good experimental animals [103].

Echinostoma echinatum (Zeder, 1803) Rudolphi, 1809

(syn. *Echinostoma lindoense* Sandground & Bonne, 1940; *Echinostoma barbosai* Jeyarasasingam et al., 1972)

This echinostome was originally described from the intestine of mammals in Germany in 1803 [104]. It has 37 collar spines arranged in 2 alternating rows [5]. This species closely resembles *Echinostoma revolutum*, and thus was once synonymized with *E. revolutum* by Diez in 1909 [104]. However, various authors acknowledge its distinctiveness [104-107]. In the meantime, *Echinostoma barbosai* Jeyarasasingam et al., 1972 and *Echinostoma lindoense* Sandground and Bonne, 1940 were synonymized with *E. echinatum* [105,108]. High prevalences (24-96%) of human infections with this echinostome were reported in 3 villages of Celebes, Indonesia, under the name of *E. lindoense* in 1940 [109]. Now this species is known to be distributed widely in Europe, Asia, and South America; natural definitive hosts can be birds and mammals [108]. Freshwater snails, including *Lymnaea*, *Planorbis*, *Anisus*, *Gyraulus*, *Biomphalaria*, and *Viviparus*, shed the cercariae [108]. Mussels, *Corbicula lindoensis*, *Corbicula sucplanta*, and *Idiopoma javanica*, contain the metacercariae [12]. *Biomphalaria glabrata* snail is another second intermediate host in Brazil [110]. Experimental definitive hosts include rats, mice, ducks, and pigeons [12,73].

Echinostoma hortense Asada, 1926

This echinostome was originally described from rats in Japan [111]. After then, it has been reported from rats in Korea [2,62, 87,88,112] and China [113]. Its head collar is armed with 27-28 collar spines arranged in 2 alteranative rows [2]. A characteristic morphology includes a laterally deviated location of the

ovary [2]. A new combination, *Isthmiophora hortensis*, was proposed for this species [114]. However, *E. hortense* should be retained until a definite conclusion could be drawn. Freshwater snails, including *Lymnaea pervia* and *Radix auricularia coreana*, shed the cercariae [2,115]. The loaches, *Misgurnus anguillicaudatus* and *Misgurnus mizolepis*, and other freshwater fish, including *Odontobutis obscura interrupta*, *Moroco oxycephalus*, *Coreoperca kawamebari*, and *Squalidus coreanus* harbor the metacercariae [2,116-118]. In Liaoning province of China, 69.7% of the loach *M. anguillicaudatus* from a market was infected with this echinostome [6]. Natural definitive hosts include rats [62,87,88,112], dogs [102], and cats [35]. Mice, rats, and humans have been shown experimentally to be susceptible to this echinostome infection [119,120]. Human infections have been reported in Japan, Korea, and China [2,5,121,122]. In Liaoning province of China, 6 out of 10 hospitalized hepatitis patients who had eaten raw loach were found infected [123]. In the Republic of Korea, an endemicity of this echinostome infection, with 22.4% prevalence, was reported among residents of Cheongsong-gun [124]. In hospitals of the Republic of Korea, clinical cases are at times diagnosed by extracting worms through gastroduodenal endoscopy [125-129].

Echinostoma ilocanum (Garrison, 1908) Odhner, 1911

This echinostome was first found from 5 prisoners in Manila, Philippines, and 21 adult flukes were recovered from 1 patient after anthelmintic treatment in 1907 [130]. Its head collar is armed with 49-51 collar spines and it characteristically has 2 deeply lobed testes [73,131]. Freshwater snails, *Gyraulus* or *Hippeutis*, shed the cercariae [6]. Large snails, *Pila conica* (Philippines) and *Viviparus javanicus* (Java), harboring the metacercariae are the source of human infections [12]. Reservoir hosts include rats and dogs [12]. Human infections have been reported from Celebes, Java, Indonesia, China, Thailand, the Philippines, and India [6,130-133]. In northern Luzon, the Philippines, the prevalence among the Ilocano population was 11% on average (7-17% by age group) [130].

Echinostoma macrorchis Ando & Ozaki, 1923

This echinostome was originally described from rats, *Rattus rattus* and *Rattus norvegicus*, in Japan [84]. The head collar is armed with 43-47 (usually 45) spines arranged in 2 alternating rows [73]. Thirty-four adult specimens were recovered from a Japanese boy in 1927 [134]. Freshwater snails, *Segmentina nitella* and *Planorbis compressus japonicus* shed the cercariae and also

harbor the metacercariae [73]. Other snails, *Cipangopaludina malleata*, *Cipangopaludina japonica*, *S. nitiella*, *Viviparus malleatus*, and the frog *Rana* sp. also harbor the metacercariae [6]. A bird, *Capella gallinago gallinago*, is another natural definitive host in Japan [135].

Echinostoma revolutum (Froelich, 1802) Looss, 1899

(syn: *Echinoparyphium paraulum* Diez, 1909)

This echinostome is the oldest species recorded in the literature [5]. It was first found in 1798 from a naturally infested wild duck *Anas boschas fereae* in Germany and reported in 1802 as *Fasciola revoluta* by Froelich [104]. Now this is the most well known echinostome species, and represents the so-called 37-collar-spined *E. revolutum* group [105]. Its geographical distribution is wide in Asia, Europe, Africa, Australia, New Zealand, and North and South America [6,104,108]. The head collar is armed with 37 spines arranged as 5-6-15-6-5, including 5 corner spines, 6 lateral spines in single rows, and 15 dorsal spines in a double row [136]. Freshwater snails, *Lymnaea* sp., *Physa* sp., *Paludina* sp., *Segmentina* sp., and *Heliosoma* sp. shed the cercariae [12]. Tadpoles, snails, or clam *Corbicula producta* harbor the metacercariae and act as the source of infection for definitive hosts [12]. In the Republic of Korea, this echinostome was reported from house rats and cats [35,75]. *Echinoparyphium paraulum*, a synonym of *E. revolutum* [12], was described from dogs in India [48] and ducks in Japan [73]. Human infection was first reported in Taiwan in 1929 [137], and the prevalence in Taiwan was once estimated to be 2.8-6.5% [6]. Human infections are also known in Yunnan and Guangdong provinces of China, Indonesia, Thailand, and Russia [12,73,138]. Ducks, geese, and muskrats are natural definitive hosts [6,104,108].

Episthmium caninum (Verma, 1935) Yamaguti, 1958

This echinostome was originally described in dogs in Calcutta, India [139]. Other *Episthmium* species have been reported from birds [48]. *E. caninum* has 24 dorsally interrupted collar spines; 12 dorsal spines are arranged in a single row and 12 lateral and ventral spines are arranged in 2 alternating rows [139]. The genus *Episthmium* was suggested to be a synonym of *Echinochasmus* [140]. Several human cases have been reported from northeast Thailand [139,141], and freshwater fish was the source of infection [141].

Fasciolopsis buski (Landkester, 1857) Odhner, 1902

This fasciolid fluke, the largest fluke among those parasitiz-

ing the human host, was first discovered in 1857 in the duodenum of an Indian man who died in London [12,142]. It is now known to be a common intestinal parasite of man and pigs in China, Taiwan, Thailand, Vietnam, Laos, Cambodia, Bangladesh, India, Indonesia, and Malaysia [4,6,142,143]. The prevalence of human infection varied according to countries, 10% in Thailand [144], 25% in Taiwan [145], 57% in China [146], and 60% in India [147]. Freshwater snails, *Segmentina* sp., *Hippeutis* sp., and *Gyraulus* sp. shed the cercariae [4,6,142]. Aquatic plants, such as, water chestnut *Eliocharis tuberosa*, water caltrop *Trapa natans*, water hyacinth *Eichhornia* sp., roots of the lotus, water bamboo *Zizania* sp., and other aquatic vegetations have the metacercariae on their body surface; metacercariae may float in the water [1,6,12]. The major mode of human infection is consumption of raw or improperly cooked aquatic plants, or peeling off the hull or skin of the plants by mouth before eating the raw nut [6]. Pigs are the main source of eggs and drainage of pig excreta in farms is an important factor for maintaining high endemicity [6].

Fischoederius elongatus (Poirier, 1883) Stiles & Goldberger, 1910

This paramphistome species is a parasite of ruminants first reported in 1883 [6]. It is infected by ingesting aquatic plants having the metacercariae [6]. Only 1 human infection was reported from Guandong, China in 1992 [6]. The patient complained of epigastric pain for several months [6].

Gastrodiscoides hominis (Lewis & McConnell, 1876) Leiper, 1913

This amphistome species was originally described from an Indian patient in 1876 [12]. Now it is known to be a common parasite of humans and pigs in India, Pakistan, Myanmar, Vietnam, the Philippines, Thailand, China, Kazakstan, Indian immigrants in Guyana, and the Volga Delta in Russia [12,142]. In human infections, worms attach to the cecum and ascending colon and may produce mucous diarrhea [12]. The planorbid freshwater snails, *Helicorbis coenosus* shed the cercariae [12], and the cercariae encyst on aquatic plants, or in tadpoles, frogs, and crayfish [6]. Pigs, mouse deers, field rats, and rhesus monkeys are reservoir hosts [6,12].

Gymnophalloides seoi Lee, Chai & Hong, 1993

This gymnophallid fluke was first discovered from a Korean woman suffering from acute pancreatitis and gastrointestinal

discomforts in 1988 [148,149]. The home village of the patient was a southwestern coastal island named Aphaedo, Shinan-gun, which was subsequently found to be a highly endemic area [150]. To date, more than 25 villages on western and southern coastal islands [151-155] and 3 non-island coastal villages [156] were identified as endemic areas. A 17th century femal mummy found in Hadong, a southern costal area of Korea, has been found to be infected with *G. seoi*; thus, this species is thought to have been prevalent in Korea for longer than several hundred years [157]. This parasite has been known to be present only in the Republic of Korea. Morphological characters of the adult parasite include a small body with a large oral sucker and a small ventral sucker, 2 short ceca, 2 compact masses of vitellaria, and a unique ventral pit [148,149]. Cercariae have not been discovered and the first intermediate host is yet unknown [149,158]. However, the oysters *Crassostrea gigas* have been proved to harbor the metacercariae [159,160]. The Palearctic oystercatchers *Haematopus ostralegus* [161] and feral cats [162] have been shown to be natural definitive hosts. Wading birds, such as, the Kentish plover *Charadrius alexandrinus*, Mongolian plover *Charadrius mongolus*, and grey plover *Pluvialis squatarola*, were experimentally proved to be susceptible to infection with this fluke [163]. Laboratory animals, including gerbils, hamsters, cats, and several strains of mice, were also found susceptible to experimental infections [164]. The metacercariae could be successfully cultured in vitro into adults using NCIC 109 medium or minimum essential medium [165,166].

Haplorchis pumilio (Looss, 1886) Looss, 1899

(syn. *Monorchotrema taihokui* Chen, 1936)

This heterophyid species, characteristically having only 1 testis and 27-39 (av. 32) minute chitinous spines on its ventro-genital sac, was originally described from birds and mammals in Egypt in 1886 [48,167]. Later the same species was discovered in Taiwan and described using the name, *Monorchotrema taihokui*, by Nishigori in 1924 [168]. Adult worms are characterized by the presence of only 1 testis and a ventro-genital-sucker complex armed with gonotyl and chitinous spines [168]. An experimental human infection with this fluke was successful by Faust and Nishigori [169], but natural human infections have been known after a report of 12 cases in 1983 in Thailand [167]. Now this fluke is known to be distributed in the Philippines, Thailand, Laos, Vietnam, South China, Taiwan, Iraque, Malaysia, India, Sri Lanka, and Egypt [6,167,170-173]. The freshwater snail, *Melania reiniana* var. *hitachiensis*, was found to shed the cer-

cariae [169,170], and freshwater fish, belonging to the Cyprinidae, Siluridae, and Cobitidae, harbor the metacercariae [170, 173]. Dogs and cats serve as the natural definitive hosts [48].

Haplorchis taichui (Nishigori, 1924) Chen, 1936

(syn. *Monorchotrema taichui* Nishigori, 1924; *Monorchotrema microrchia* Katsuda, 1932; *Haplorchis microrchis* Yamaguti, 1958)

This heterophyid species, characteristically having only 1 testis and 14-20 (av. 15) large chitinous spines on its ventro-genital sac, was originally described from birds and mammals caught in Taiwan [168,169,174]. An experimental human infection with this fluke was successful by Faust and Nishigori [169], but natural human infections have been known after reports in the Philippines [175]. Now it is known to be distributed widely in Taiwan, the Philippines, Bangladesh, India, Sri Lanka, Palestine, Iraque, Egypt, Malaysia, Thailand, Laos, Vietnam, and South China [4,6,170-172,175-177]. Freshwater snails, *Melania oblique-granosa*, *Melania juncea*, or *Melanoides tuberculata*, serve as the first intermediate hosts [169,170]. Freshwater fish, including *Cyclocheilichthys repasson*, *Cyprinus auratus*, *Cyprinus carpio*, *Gambusia affinis*, *Hampala dispar*, *Labiobarbus leptocheila*, *Puntius binotatus*, *Puntius brevis*, *Puntius gonionotus*, *Puntius leicanthus*, *Puntius orphoides*, *Puntius palata*, *Pseudorasbora parva*, *Rhodeus ocellatus*, and *Zacco platypus* [170,173] and *Raiamas guttatus*, *Mystacoleucus marginatus*, and *Henichoryhnchus siamensis* [178] harbor the metacercariae. Dogs, cats, and birds are the natural definitive hosts [48].

Haplorchis vanissimus Africa, 1938

This heterophyid species was originally described from a naturally infected man in the Philippines [175]. Thereafter, this fluke was reported from pelicans and wild mammals in Australia [179]. Freshwater fish serve as the second intermediate host [6].

Haplorchis yokogawai (Katsuta, 1932) Chen, 1936

(syn. *Monorchotrema yokogawai* Katsuda, 1932)

This heterophyid species, characteristically having only 1 testis and numerous (uncountable) minute chitinous spines on its ventro-genital sac, was originally described from dogs and cats experimentally fed the mullet *Mugil cephalus* in Taiwan [180]. An experimental human infection with this fluke was successful by Katsuda [180], but natural human infections have been known after reports in the Philippines [175]. Now this fluke is known to occur in the Philippines, South China, Malaysia, Indonesia, Thailand, Laos, Vietnam, India, Australia, and Egypt as human or animal infections [6,170,172,176]. Freshwater snails,

Melanoides tuberculata or *Stenomelania newcombi*, shed the cercariae [170]. Freshwater fish, including *Cylocheilichthys armatus*, *Hampala dispar*, *Labiobarbus leptocheila*, *Misgurnus* sp., *Mugil* spp., *Onychostoma elongatum*, *Ophicephalus striatus*, and *Puntius* spp. harbor the metacercariae [170,173]. Dogs, cats, cattle, and other mammals are natural definitive hosts [48].

Heterophyes nocens Onji & Nishio, 1916

(syn. *Heterophyes katsuradai* Ozaki & Asada, 1926)

This heterophyid species, which characteristically has 50-62 chitinous spines on its gonotyl, was first reported from experimental dogs and cats fed the mullet *Mugil cephalus* in Japan [181]. It is now known to occur as human infections in Japan, China, and the Republic of Korea [182,183-187]. Brackish water snails, *Cerithidea cingulata* (= *Tympanotonus microptera*), shed the cercariae [4,40]. Brackish water fish, such as the mullet *Mugil cephalus* or goby *Acanthogobius flavimanus*, harbor the metacercariae [2,187]. Domestic or feral cats are naturally infected with this fluke [35,162,188]. In the Republic of Korea, prevalences ranging from 10% to 70% were detected among residents in southwestern coastal areas, including islands [2,189]. In Japan, human *H. nocens* infections were reported from Kochi, Chiba, Yamaguchi, Chugoku, Hiroshima, and Shizuoka Prefectures [190,191].

Heterophyopsis continua (Onji & Nishio, 1916) Yamaguti, 1958

(syn. *Heterophyes continuus* Onji & Nishio, 1916; *Heterophyes expectans* Africa & Garcia, 1935)

This heterophyid species, characteristically having an elongate body, a genital sucker located separately from the ventral sucker, and 2 obliquely tandem testes, was first described from experimental cats fed the mullet *Mugil cephalus* that harbored the metacercariae in Japan [2,181]. Unknowing this species, Africa and Garcia [192] described *Heterophyes expentans* as a new species from dogs, which later has been synonymized with *H. continua* [48]. Human infections were first mentioned in Japan, without details [193]. After 2 natural human infections were first discovered [101], 8 more human cases have been confirmed in the Republic of Korea [94,151,152,194]. The snail host is unknown. Brackish water fish, including the perch *Lateolabrax japonicus*, goby *Acanthogobius flavimanus*, and shad, *Clupanodon punctatus* [195,196], conger eel *Conger myriaster* [197], and sweetfish *Plecoglossus altivelis* [198] harbor the metacercariae. Cats [35, 188], ducks [181], and sea-gulls [8] are the natural definitive

hosts. Cats [181], dogs [101,195] and domestic chicks [199,200] are experimental host in the laboratory.

Hypoderaeum conoideum (Block, 1872) Diez, 1909

This echinostome, having a small inconspicuous head collar with 47-53 (usually 49) collar spines arranged in 2 alternating rows, was discovered in 1872 from birds and now is known to be an intestinal fluke of the duck, goose, and fowl in Europe, Japan and Siberia [48,73]. Freshwater snails, *Planorbis corneus*, *Indoplanorbis exustus*, *Lymnaea stagnalis*, *Lymnaea limosa*, *Lymnaea ovata*, and *Lymnaea rubiginosa*, shed the cercariae, and snails and tadpoles are the second intermediate hosts [48,201]. *Lymnaea tumida* in Russia [202] and *Lymnaea peregra* and *Lymnaea corvus* in Spain [203] were experimentally proven to be potential first intermediate hosts. In Thailand, this echinostome was reported from humans and birds [201,204]. In an area of northeast Thailand, 55% of 254 residents were found infected with this echinostome [204].

Isthmiophora melis (Schrank, 1788) Lühe, 1909

(syn. *Euparyphium melis* Railliet, 1919, *Euparyphium jassyense* Leon & Ciurea, 1922)

This fluke, characteristically having 27 collar spines, 19 of which are arranged in uninterrupted double rows and 4 are corner spines on each side, was described from rodents and carnivores in Europe and North America [48,73]. Its taxonomic position was unstable, either placing it in *Euparyphium* [12] or *Echinostoma* [201]. However, recently it has been assigned to the genus *Isthmiophora* [114]. Human infection with *I. melis* was first confirmed by Leon in 1916 in a diarrheic patient in Rumania and then at autopsy of a Chinese patient [12]. Further human cases were reported in China and Taiwan [6]. The snail *Stagnicola emarginata angulata* is the first intermediate host in the region of Douglas Lake, Michigan, and tadpoles serve as the second intermediate host [12]. The source of human infections in China was presumed to be the loach, a kind of freshwater fish [6]. Domestic and wild animals are natural definitive hosts in Russia [205].

Metagonimus minutus Katsuta, 1932

This heterophyid species, characterized by smaller sized uterine eggs, was reported from experimental mice and cats fed mullets harboring the metacercariae in Taiwan [206]. This parasite is listed among the human-infecting intestinal trematodes [6,12], but no literature background is traceable [4].

***Metagonimus miyatai* Saito, Chai, Kim, Lee & Rim, 1997**

This heterophyid fluke, having widely separated 2 testes with 1 testis located near the posterior end of the body, was first found in Japan by Miyata in 1941, but its taxonomic position was not established until it was reported as a new species in 1997 [207]. This fluke is morphologically different from *M. yokogawai* and *M. takahashii* in the position of the posterior testis (separated considerably from the anterior one), the distribution of vitelline follicles (never crossing over the posterior testis), and the intermediate size of eggs (28-32 μm) [207]. Human infections with this species were first reported in 1980 (under the name of *Metagonimus* sp.) from people along the Gum River, the Republic of Korea [208]. Subsequently, high prevalences were reported among people residing around the Daechong Reservoir and its upper reaches [209], the upper reaches of the Namhan River [210], the Hantaan River basin [211], and the western inland of Gangwon-do [212]. In Japan, small rivers of Shizuoka Prefecture, Japan, were found to be endemic areas of this fluke [213]. Freshwater snails, *Semisulcospira globus* [209], *Semisulcospira libertina*, or *Semisulcospira dolorosa* [214], are the first intermediate host. Freshwater fish, including the sweetfish, dace, common fat-minnow *Morocco steindachneri*, pale chub *Zacco platypus*, and dark chub *Zacco temminckii*, harbor the metacercariae [207]. Mice, rats, hamsters, and dogs are experimental definitive hosts [2,215]. Natural definitive hosts are unknown.

***Metagonimus takahashii* Suzuki, 1930**

This heterophyid fluke, having slightly separated 2 testes and extensive distribution of vitellaria near the posterior end of the body, was first reported in Japan from mice and dogs fed freshwater fish other than the sweetfish [216]. This species has been found also in the Republic of Korea, particularly along small streams in various inland areas [2]. Morphologically, it differs from *M. yokogawai* in the position of the 2 testes, the distribution of vitelline follicles, and the egg size (*M. yokogawai*, 28-30 μm ; *M. takahashii*, 32-36 μm) [2,4]. The Koga type of *Metagonimus* encysting in the dace *Tribolodon* spp. [217] is regarded as a synonym of *M. takahashii* [4,218]. The presence of this species in the Republic of Korea was first documented by recovery of adult flukes from experimental rabbits in 1960 [219]. With regard to human infections, adult flukes were first confirmed in 1993 in inhabitants of Umsong-gun, Chunchungnam-do, along the upper reaches of the Namhan River [4,210]. The inhabitants were mixed-infected with *M. miyatai*, with an egg positive rate of 9.7% for both species [210]. Freshwater snails, *Semisulcospira*

coreana or *Koreanomelania nodifila* [220], may be first intermediate host. Freshwater fish, including the crussian carp *C. carassius* [219], carp *C. carpio* [217], dace *Tribolodon taczanowskii* [218], and perch *Lateolabrax japonicus* [221], harbor the metacercariae. Different strains of mice could be experimentally infected with fluke [215]. There are no reports on reservoir hosts.

***Metagonimus yokogawai* (Katsurada, 1912) Katsurada, 1912**

This heterophyid species, morphologically characterized by closely adjacent 2 testes and moderate distribution of vitellaria, is the most common intestinal fluke infecting humans in the Far East [2,4]. This fluke was first found from humans and experimental dogs in Taiwan by S. Yokogawa in 1911 [40]. It was originally named as *Heterophyes yokogawai* Katsurada, 1912 and then changed into *Metagonimus yokogawai* in the same year [40]. Human infections were reported from Japan, Korea, Siberia, Israel, the Balkan states, and Spain [6]. In the Republic of Korea, almost all large and small streams in eastern and southern coastal areas are endemic foci of metagonimiasis [4,222,223]. The Sumjin, Tamjin, and Boseong Rivers, Geoje Island, and Osip Stream (Gangwon-do) were the highest endemic areas with 20-70% egg positive rates in the villagers [2,222,224,225]. The high prevalence of *M. yokogawai* along the Tamjin River basin is persisting, although the intensity of infection among the villagers has been reduced [226]. Human infections have also been found in Guangdong, Anhui, Hubei, and Zhejiang provinces of China and Taiwan [6,183]. In Japan, the prevalence rate became negligible after the 1970s, except in some areas surrounding the Hamana Lake [227]. In fish hosts, however, small rivers of Shizuoka prefecture were prevalent with *M. yokogawai* metacercariae [213]. In Russia, the endemic areas include the Amur and Ussuri valleys of Khabarovsk territory where the prevalence in ethnic minority groups was between 20% and 70% [6]. In the north of Sakhalin Island, the infection rate was 1.5% in Russians and 10% in ethnic minorities. Sporadic cases were also reported in Amur district and Primorye territory [6]. Freshwater snails, *Semisulcospira coreana* or *Semisulcospira libertina*, shed the cercariae [220]. The sweetfish *Plecoglossus altivelis* [2], the dace *Tribolodon* sp. [218], and the perch *Lateolabrax japonicus* [228] serve as second intermediate hosts. Dogs, rats, and cats are the natural definitive hosts [88,102,229]. Variable mouse strains were found more susceptible to *M. yokogawai* infection and less susceptible to *M. miyatai* and *M. takahashii* infections [215].

***Neodiplostomum seoulense* (Seo, Rim & Lee, 1964) Hong and Shoop, 1995**

This neodiplostomum species was originally described from naturally infected house rats in the Republic of Korea [87]. It is morphologically unique due to its bisegmented body, presence of a tribocytic organ, 2 butterfly-shaped testes, and a wide distribution of vitellaria in the anterior body to the level of the ventral sucker [230]. This parasite is now known to be distributed countrywide among rodents in the Republic of Korea, but predominantly in mountainous areas [4,230]. This fluke was reported to be present also in a northeastern part of China [231]. The first human case was found in 1982 in a young man suffering from acute abdominal pain and fever, who had a history of consuming improperly cooked snakes 7 days prior to admission to a hospital [232]. Since then, 26 more human cases were found among soldiers who had eaten raw snakes during their survival training [233-235]. Freshwater snails, *Hippeutis cantori* [236] and *Segmentina (Polypylis) hemisphaerula* [237] shed the cercariae. Tadpoles and frogs of *Rana* sp. are the second intermediate host [236], and the snake *Rhabdophis tigrina* is regarded as a paratenic host [230]. Mice, rats, and guinea pigs have been found to be susceptible laboratory hosts [230]. Rats, *Rattus norvegicus* [88, 236] and mice *Apodemus agrarius* [238] are the natural definitive hosts.

***Phaneropsolus bonnei* Lie Kian Joe, 1951**

This lecithodendriid species, which characteristically has 8 vitelline follicles on each anterolateral side, was first described from a human autopsy in 1951 in Jakarta, Indonesia, and then found in monkeys in Malaysia and India in 1962 [239]. After then, this fluke was reported from 15 human autopsies in Udornthani Provincial Hospital, northeast Thailand [240,241]. Subsequently, high prevalences of this fluke infection were found in northeastern provinces of Thailand [14]. Also in Lao PDR, a total of 1,303 adult specimens were recovered from 27 people residing in Mekong riverside areas of Savannakhet, Khammouane, and Saravane Provinces [62,171,176]. The egg morphology is very similar to that of heterophyid fluke or *Opisthorchis viverrini* [242,243]. Metacercariae were discovered in naiads and adult dragon- and damselflies in Thailand [244]. Local people in northeast Thailand and Laos sometimes eat naiads of these insects [171,176,244].

***Phaneropsolus spinicirrus* Kaewkes et al., 1991**

This lecithodendriid species was described as a new species

from a human infection case in northeast Thailand [245]. No further information is available on this parasite.

***Plagiorchis harinasutai* Radomyos, Bunnag & Harinasuta, 1989**

This plagiorchiid species was reported as a new species in 1989 based on worms recovered from 4 human cases [246]. No other information is available on this fluke.

***Plagiorchis javensis* Sandground, 1940**

This plagiorchiid species was originally described from a human case in Indonesia [247]. Later 2 more cases were found in Indonesia [6]. Larval insects are presumed to be the source of infection, and birds and bats are reservoir hosts [6].

***Plagiorchis muris* (Tanabe, 1922) Shul'ts & Skvortsov, 1931**

This plagiorchiid species, which has a laterally located ovary, 2 tandem testes, and extensive distribution of vitellaria from the posterior end to the level of the pharynx, was originally described in Japan based on worms recovered from mice experimentally infected with the metacercariae [248]. An experimental human infection was reported in the U.S. [249]. Natural human infections have been reported in Japan [250] and the Republic of Korea [251]. Freshwater snails, *Lymnaea pervia* in Japan [248] and *Stagnicola emarginata angulata* in the U.S. [249], shed the cercariae. Aquatic insects (mosquito larvae), insect naiads, fresh water snails, and fresh water fish [248-251] harbor the metacercariae. Rats are an experimental definitive host [251]. House rats in Japan [248], and rats [87,88], mice [252], and cats [35] in the Republic of Korea are the natural definitive hosts.

***Plagiorchis philippinensis* Sandground, 1940**

This plagiorchiid fluke was first recovered at autopsy of a resident in Manila, Philippines by Africa and Garcia in 1937 [6, 247]. Infection was acquired by eating insect larvae [6]. Birds and rats are reservoir hosts [6].

***Plagiorchis vespertilionis* (Müller, 1780) Braun, 1900**

This plagiorchiid species was originally described from the brown long-eared bat in Europe, and then found in many countries, including the Republic of Korea [253]. This fluke has been found infected in a human in the Republic of Korea [254]. The patient had habitually eaten raw flesh of snakehead mullet and gobies [254]; however, it is uncertain whether these fish took the role for a source of infection. Its snail host is *Lymnaea*

stagnalis, and the second hosts reported include mosquito larvae, caddis-fly larvae, mayfly larvae, and dragonfly nymphs [255].

Procerovum calderoni (Africa and Garcia, 1935) Price, 1940

This heterophyid species was first reported from 5 dogs and a native Filipino [256]. Later, it was reported from China and Africa [201]. Brackish water snails, *Thiara riquetti*, shed the cercariae [257]. Freshwater fish, *Ophiocephalus striatus*, *Glossogobius giurus*, *Mollienesia latipinna*, *Mugil* sp., and *Creisson validus* harbor the metacercariae [6,257,258].

Procerovum varium Onji & Nishio, 1916

This heterophyid parasite was described from experimentally infected dogs with the metacercariae in the mullet *Mugil cephalus* in Japan [181]. Experimental human infections were reported successful [259], but there have been no reports on natural human infections. It is now known to be distributed in China, the Philippines, Australia, India [260], and the Republic of Korea [35]. Cats are the natural definitive hosts [35].

Prosthodendrium molenkampii Lie Kian Joe, 1951

This lecitodendriid species, which characteristically has 12-30 vitelline follicles on each anterolateral side, was first described from 2 human autopsies by Lie Kian Joe in 1951 in Jakarta, Indonesia [244]. After then, this fluke was found in 14 human autopsies in Udomthani Provincial Hospital, Thailand [241,244]. Later, high prevalences were reported in different areas of northeast Thailand [14]. In Lao PDR, a total of 8,899 adult specimens were recovered from 52 infected people residing along the Mekong riverside areas of Vientian Municipality, Savannakhet, Khammoune, and Saravane Provinces [62,171,176]. Metacercariae were discovered in naiads and adult dragon- and damselflies in Thailand [244]. Local people in northeast Thailand and Laos at times eat naiads of these insects [62,171,244].

Psilorchis hominis Kifune & Takao, 1973

This psilostomid fluke was described from a 48-year-old Japanese patient, who had a mixed infection with *E. macrorchis*, after anthelmintic medication [261]. No other information is available on this species.

Pygidiopsis summa Onji & Nishio, 1916

This heterophyid species, characterized by a small concave

body, median location of the ventral sucker, unique morphology of the ventrogenital apparatus, and side-by-side location of the two testes [262], was first found in dogs fed brackish water fish infected with the metacercariae in Japan [181], and then reported from the Republic of Korea [2]. Human infections were reported by detection of eggs in the feces in 1929 [263] and of adult flukes in 1965 in Japan [182]. In the Republic of Korea, 8 human infections were discovered from a salt-farm village of Okku-gun, Chollabuk-do [184]. In 2 other coastal areas of the Republic of Korea, 18 and 5 heterophyid egg-positive people were found to be infected with this fluke [151,152]. It is now known to be distributed widely along the western and southern coastal islands of the Republic of Korea [188]. Brackish water snails, *Cerithidea* sp. or *Tympanotonus* sp., shed the cercariae [4]. Brackish water fish, including the mullet *Mugil cephalus* and goby *Acanthogobius flavimanus*, harbor the metacercariae [187,196,264,265]. Domestic or feral cats are the natural definitive hosts [35,162,188].

Spelotrema brevicacaeca (Africa & Garcia, 1935) Tubangui & Africa, 1939

This microphallid fluke was originally reported under the name *Heterophyes brevicacaeca* from a human autopsy case [256], and then 11 cases in the Philippines [175]. Eggs of this fluke were found in the heart, brain, and spinal cord of the patients who died of acute cardiac dilatation [175,256]. Brackish water crabs, *Carcinus maenas*, and shrimps, *Macrobrachium* sp., harbor the metacercariae in the Philippines [12].

Stellantchasmus falcatus Onji & Nishio, 1916

This heterophyid species, characterized by the slightly deviated ventral sucker to the right side of the body and the presence of an elongated sac-like seminal vesicle on the opposite side of the ventral sucker [2], was first described from cats experimentally fed the mullet harboring the metacercariae in Japan [181]. First human infections [263] and successive cases [40,266] were reported in Japan. Thereafter, human infections have been reported in many Asian-Pacific countries; the Philippines, Hawaii, Japan, Palestine, Thailand, Vietnam, and the Republic of Korea [2,4,48,172,267,268]. Brackish water snails, *Stenomelania newcombi* or *Thiara granifera*, shed the cercariae [42,269]. The mullet [2,270] and half-beaked fish *Dermogenus pusillus* [271,272] harbor the metacercariae. Rats were the better experimental hosts than mice [273]. Cats are the natural definitive hosts [35,263].

***Stellantchasmus formosanus* Katsuta, 1931**

This heterophyid species was originally described from experimentally infected cats, dogs and mice with the metacercariae encysted in the mullet *Mugil cephalus* in Taiwan [274]. An experimental human infection was reported in Taiwan [274]. No reports have been available on natural human infections.

***Stellantchasmus pseudocirratu*s (Witenberg, 1929) Yamaguti, 1958**

(syn. *Stellantchasmus amplicaealis* Katsuta, 1932)

This heterophyid species was first described from naturally infected dogs and cats in Palestine [275], followed by cats, dogs, and mice fed mullet in Taiwan [276]. Human infections were reported in the Philippines and Hawaii [48,175]. The mullet, *Mugil* sp., is the second intermediate host [48,275].

***Stictodora fuscata* (Onji and Nishio, 1916) Yamaguti, 1958**

This heterophyid species, morphologically characterized by the presence of a gonotyl, which is superimposed on the ventral sucker and armed with 12 chitinous spines, a prominent metraterm, and 2 testes located obliquely in the middle field of the body, was originally described from cats experimentally fed infected mullets in Japan [4,181]. The first human infection with this fluke, under the name *Stictodora* sp., was found in a young Korean man, who regularly ate raw mullets and gobies [277]. Subsequently, 13 additional human cases were detected in a seashore village in the southwestern coastal area [2,4]. The gobies, *Acanthogobius flavimanus*, are the second intermediate host [196,278]. Cats (*Felis catus*) were used as an experimental definitive host [196]. Feral cats were found naturally infected with this fluke [35].

***Stictodora lari* Yamaguti, 1939**

This heterophyid fluke, morphologically characterized by a gonotyl armed with 70-80 min spines, was first found in the small intestine of the sea gull *Larus crassirostris* in Japan [8,279]. Its adult flukes were first detected in 6 Korean people who resided in 2 southern coastal villages [280]. Brackish water gastropods, *Velacumantus australis*, shed the cercariae in Australia [281]. A number of estuarine fish, including the goby, *Acanthogobius flavimanus*, harbor the metacercariae [278,281]. Cats and dogs were used as experimental definitive hosts [278]. Feral cats are the natural definitive hosts [35].

MOLECULAR STUDIES**Echinostomes**

There has been no full genome sequencing project for echinostomes, and only 474 DNA sequences are available in Genbank, as of January 2008 [282]. Most of these DNA data have been generated for phylogenetic and taxonomic studies. This low number of genome sequences in the database often makes it fail to identify the protein from peptides obtained from echinostomes [282]. *Echinostoma paraensei* is the most frequently studied species, and 358 of 474 reported sequences were obtained from the sporocysts RNA of this species [283]. *E. revolutum* and *E. caproni* are the next frequently studied species. The rest of the deposited sequences mainly represent molecules used in taxonomic and phylogenetic studies, including ribosomal or mitochondrial DNA [282]. There have been no studies on microarrays or reverse genetics of echinostomes.

Most of the ribosomal molecules sequenced corresponded to the internal transcribed spacers (ITS1 and ITS2), and 5.8S, 18S, and 28S genes [282]. With respect to mitochondrial markers, those studied included NADH dehydrogenase subunit 1 (ND1) and the cytochrome oxidase 1 [282]. Morgan and Blair [284] used ITS1, ITS2, and 5.8S ribosomal DNA to study 8 echinostome species. They found that 5 ('*revolutum*' group) of the 8 species revealed distinguishable ITS data from the others, but within the '*revolutum*' group, they found only low sequence divergence. Later, the same authors [285] reported that ND1 is a more suitable marker for species and strain detection of echinostomes within the group. However, new cDNA and genomic libraries are required for further echinostome genetic studies.

The study of echinostome proteins has been initiated in 1968 by Taft and Fried [286], who first showed an indirect evidence of the presence of cytochrome oxidase in *E. revolutum*. Subsequently, larval antigens of *E. lindoense* were obtained from the snail host, *Biomphalaria glabrata* [287]. The cytochrome c oxidase of *E. trivolvis* was successfully purified by Fujino et al. [288]. Since recently, studies on proteomics using 2-dimensional electrophoresis (2DE) have been performed. Using 2DE, various kinds of proteins, including actin, tropomyosin, and paramyosin, glycolytic enzymes, including endolase, glyceraldehydes 3P dehydrogenase (GAPDH), and aldolase, detoxifying enzymes, in particular, glutathion-S-transferase, and the stress-related protein, Hsp 70 have been identified from *E. friedi* [289]. The major limitation in proteomic studies on echinostomes is the lack of sufficient genomic data.

Fasciolopsis buski

Only a few molecular studies have been performed on *F. buski*. Two sequences corresponding to the 18S rRNA gene are deposited in Genbank [142]. Using this DNA sequence, fluke specimens vomited by a child was confirmed as *F. buski* [290]. Species-specific ITS sequences of ribosomal DNA were described, and the sequences of eggs and adults were identical in length and composition [291]. Phylogenetically, *F. buski* resembles closely the other members of the family Fasciolidae, including *Fasciola hepatica* and *F. gigantica* [291].

Gymnophalloides seoi

Few studies have been performed on the molecular biology, genetics, and proteomics of *G. seoi*. Crude extracts of *G. seoi* containing antigenic proteins were found to enhance the expression of Toll-like receptors 2 (TLR2) and 4 (TLR4) mRNAs on HT29 cells, a human intestinal cell line, cultured in vitro [292]. However, monoclonal antibodies against *G. seoi* 46 kDa antigen revealed no significant effects on the TLR2 and TLR4 mRNA expression on HT29 cells [292], therefore, the responsible antigenic proteins should be further clarified.

With reference to other gymnophallid species, 10 polymorphic microsatellite loci were isolated and characterized for a *Gymnophallus* species [293]. The loci included G1C10, G3A4, GYM6b, GYM8, GYM9, GYM11, GYM12, GYM14, GYM15, and GYM16, and in each locus, 3-23 alleles were detected in 24 metacercariae isolated from the bivalve host [293]. These results show that there are marked heterozygosities among the metacercariae of this gymnophallid species. Subsequently, 406 metacercariae of the same species isolated from 15 cockles were genotyped using 6 of the 10 microsatellite loci, and as many as 400 unique genotypes were identified [294]. The results supported the role of the second intermediate host as an accumulator of genetic diversity in the trematode life cycle [294].

Proteomics are poorly known in *G. seoi*. In 2 papers, cysteine proteinases were isolated from the metacercariae and adults, and functionally characterized [295,296].

Metagonimus and other heterophyids

A SDS-PAGE/immunoblot analysis of crude extracts of *M. yokogawai* metacercariae showed that out of 14 protein bands found, 11 positively reacted with infected human sera, and among them 66 kDa and 22 kDa proteins were the most specific antigens [297].

Three species of *Metagonimus*, i.e., *M. yokogawai*, *M. takahashii*,

and *M. miyatai*, can be genetically differentiated using PCR-RFLP analysis, where ITS1 site of ribosomal RNA and mitochondrial cytochrome c oxidase I (mtCOI) gene were targeted [298-300]. Especially in mtCOI gene, restriction patterns with Rsa I and Alu I produced differentially fragmented banding patterns [298]. PCR-based random amplification of polymorphic DNA (RAPD) technique using random 10-mer oligonucleotide primers also showed distinguishable banding patterns between *M. yokogawai* and *M. miyatai* [299].

Studies on the number of chromosomes and karyotypes of the genus *Metagonimus* supported strong evidence of genetic differences between *M. yokogawai* and *M. miyatai* [301]. They showed that the number of bivalents in the first meiotic division of *M. takahashii* was 9 ($n = 9$) and the diploid number of *M. miyatai* and *M. yokogawai* were 18 ($2n = 18$) and 32 ($2n = 32$). In addition, chromosomes of *M. miyatai* consisted of 1 pair of metacentric, 7 pairs of submetacentric, and 1 pair of telocentric chromosomes, whereas those of *M. yokogawai* consisted of 2 pairs of metacentric, 11 pairs of submetacentric, and 3 pairs of subtelocentric chromosomes [301]. Similarly, simple sequence repeat anchored PCR using primers of 3' or 5' termini of the (CA) n repeats showed distinguishable banding patterns among the 3 species of *Metagonimus*, suggesting that they have different genotypes [302].

The sequence comparison of 28S ribosomal DNA and mtCOI of 3 *Metagonimus* species also supported *M. miyatai* as a separate species [303]. The length of the rDNA sequence in 28S D1 region of the 3 *Metagonimus* species was 248 bp and its G+C content was 52% [303]. The length of the mtCOI sequence averaged 400 bp with a G+C content ranged from 44% in *M. miyatai* and 46% in *M. takahashii* to 47% in *M. yokogawai* [303]. Nucleotide sequence differences between species were 23.0% (92/400 bp) between *M. miyatai* and *M. takahashii*, 16.2% (65/400 bp) between *M. miyatai* and *M. yokogawai*, 13.2% (53/400 bp) between *M. takahashii* and *M. yokogawai* [303]. *M. takahashii* and *M. yokogawai* are placed in the same clade supported by the sequence and phylogenetic tree analysis of 28S D1 rDNA and mtCOI gene by neighbor-joining and parsimony method; *M. miyatai* is placed in a different clade from 2 other *Metagonimus* species [303].

With regard to other heterophyid species, only a few studies have been performed on the molecular biology, genetics, and proteomics. Sequencing and aligning of the 18S (SSU) rDNA of 6 heterophyid species, including *H. taichui*, *H. pumilio*, *Centrocestus* sp., *Pygidioopsis genata*, *Phagicola longa*, and *Dexiogonimus*

ciureanus, revealed interspecific variations [304]. However, due to high sequence similarities, it was necessary to perform PCR-RFLP of the entire 18S subunit to differentiate between *Haplorchis* and *Dexiagonimus* [304]. A high annealing temperature (HAT)-based random amplified polymorphic DNA (RAPD) technique has been developed to detect DNA of *H. taichui* and *S. falcatus* [305,306].

Neodiplostomum seoulense

Few studies have been done regarding the genetic characteristics of *N. seoulense*. Recently, in order to support data on phylogeny of 3 *Neodiplostomum* species distributing in the Republic of Korea, i.e., *N. seoulense* [230], *N. leei* [307], and *N. boryongense* [308], mitochondrial COI gene was analyzed, and positive results were obtained (to be published).

A cysteine protease, with the molecular weight of 54 kDa, was purified from the crude extract of *N. seoulense* adults, although its function was suggested to aid nutrient uptake, rather than host tissue lysis [309]. Two cystatin-binding cysteine proteinases were successfully purified from *N. seoulense*, and they were found useful for cystatin-capture ELISA of human *N. seoulense* infection [310]. Other proteases and unique proteins should be purified and characterized to better understand host-parasite relationships.

PATHOGENICITY

Echinostomes

The pathogenicity of echinostomes is complex and diverse depending on a wide variety of parasite- and host-side factors [5,6,311]. Mechanical irritation by the flukes and allergic reactions due to their toxic metabolites constitute the 2 important elements responsible for the pathogenesis. In this respect, the intensity of infection, in terms of individual worm load, may be highly important in the pathogenicity and severity of the disease [5,73,105]. Fried et al. [312] compared the host pathologies between single and multiple infections with an animal echinostome, *E. caproni*, in hamsters. Certain species of echinostome seem to be more pathogenic to humans, since a fatal human case of *A. malayanum* (under the name *A. mehrai*) was reported in India, from whom several hundred worms were recovered at autopsy [5,313]. Bowel perforation due to *A. malayanum* (under the name *A. mehrai*) infection was also reported in an Indian child [314].

In echinostome-infected rodent hosts, intestinal histopathol-

ogy was studied in *E. revolutum* [315,316], *E. hortense* [2,317], *E. caproni* [311,318], and *E. trivolvis* infection [319]. Adult worms were located in the lumen of the upper small intestine of the rodents and pathological changes occurred chiefly near the attachment sites of the echinostomes [2,311]. The flukes were tightly pinching and sucking the mucosa with their oral suckers, and it seemed that the mucosal layer was eventually eaten by the flukes [5]. Villous atrophy, crypt hyperplasia, inflammation of the stroma, and decreased villus/crypt ratios were observed in the small intestine of experimentally infected animals [316,317]. In focal areas, massive destruction and detachment of the villi, at times complete loss of the mucosal integrity and ulcerations, were observed [2,316]. These features are similar to those observed in severe catarrhal enteritis [5]. In pigs naturally infected with *A. oraoni*, fatal diarrhea was developed, and autopsy revealed massive parasite infections with hemorrhagic and edematous mucosa of the jejunum and duodenum [27].

Fasciolopsis buski

The severity of the disease is markedly variable depending on parasite load [142]. Adult flukes can cause mechanical damage to the intestinal mucosa and extensive intestinal and duodenal erosions, ulceration, hemorrhages, abscess, and catarrhal inflammation may occur [1,320]. Absorption of toxic and allergic worm metabolites can cause ascites and facial, orbital, and generalized edema [142]. A case of heavy *F. buski* infection that caused intestinal perforation has been reported [321].

Gymnophalloides seoi

The pathogenicity of *G. seoi* is not well understood partly because no suitable experimental host has been found. It was observed in mice, though they are not a good host, that *G. seoi* parasitizes the small intestine, chiefly the duodenum and jejunum, pinching and sucking the intestinal villi with their oral suckers [322]. The infected intestinal mucosa showed villous atrophy and crypt hyperplasia, with inflammatory reactions in the villous stroma and the crypt [4,322]. The histopathological changes were generally not so severe, and the mucosal integrity was restored around day 14-21 post-infection [322]. In immunocompetent mice, the worms did not invade the submucosal layer; however, in immunosuppressed mice, worms invaded the submucosa [322]. In a colon cancer patient who received anti-cancer chemotherapy, a *G. seoi* worm was found to have penetrated into the colonic lymphoid tissue [323]. Mechanical irritation and antigenic stimulation by the flukes are considered to

be important in the pathogenicity [4,322].

The small intestine is presumed to be the normal habitat of *G. seoi* in the human host [4]. In experimental rodents this was evidenced [164,322]. However, acute pancreatitis was diagnosed in 1 patient [148], and diabetes mellitus was accompanied in 2 patients [324]. Therefore, *G. seoi* is suspected to invade the pancreatic duct in humans [4]. It is of note that other gymnophalids were found in the bursa Fabricii and gallbladder of shorebirds as well as in the intestine [48]. Experiments using larger animals, such as monkeys, seem to be necessary to verify this possibility [4]. *G. seoi* was found to be able to invade the submucosa of immunosuppressed mice [322]. Therefore, there is a possibility that the eggs may be transferred to remote organs to cause erratic parasitisms in immunocompromised hosts, as reported in heterophyid infections [175].

Metagonimus and other heterophyids

Intestinal histopathology was studied in *M. yokogawai* [325-327], *P. summa* [328], and *C. armatus* [329] using experimental animals, including mice, rats, cats, and dogs. Adult flukes of *M. yokogawai* were found to parasitize the middle part of the small intestines, and invade the crypt of Lieberkühn in the early stage of infection (by day 2-3 post-infection), and localize between villi in later stages [4,325-327].

In *M. yokogawai* infection, the major histopathological findings included villous atrophy and crypt hyperplasia, with variable degrees of inflammatory reactions [4,325]. The infected mucosa shows blunting and fusion of the villi, edema of the villus tips, congestion and inflammatory cell infiltrations in the villous stroma, and decreased villus/crypt height ratios [2,4]. Very similar intestinal histopathology was reported in a naturally infected human with *M. yokogawai* [330]. The location of worms was confined to the intestinal mucosa in immunocompetent hosts [327,331,332]; however, immunosuppression allowed a deeper invasion of the worms into the submucosa in mice [333]. Immunosuppression also enhanced the survival of worms and prolonged the life spans in the same mouse strain [333,334]. Decreased enzyme activities were observed in acute infections with *M. yokogawai* [335]. In dogs experimentally infected with *M. yokogawai*, it was suggested that watery content might be a result of a poor absorption of the intestinal secretions from the secretory crypt cells [336].

In *M. miyatai*-infected mice, similar intestinal histopathology was observed, although the degree of mucosal damage was less severe than in *M. yokogawai*-infected mice, as represented by

stronger expression patterns of the proliferating cell nuclear antigen (PCNA) in the intestinal mucosa [337]. In *P. summa*-infected mice, similar features were also observed; the worms caused severe villous atrophy and crypt hyperplasia, with inflammation of the villous stroma [328]. In experimental *C. armatus* infection in rats, the worms caused mechanical irritation and mucosal inflammations in the small intestines from as early as 3 days after the infection [329].

With respect to the pathogenicity of heterophyid flukes, it should be noted that several species, namely *S. falcatus*, *Haplorchis* spp., and *Procerovum* spp., can cause erratic parasitism in man, which is often fatal [175]. The 3 most frequently affected sites were the heart valve, brain, and spinal cord, where eggs and adult flukes originating from the intestinal mucosa embolized in the blood vessels [175]. Eggs of *H. nocens* (under the name *H. heterophyes*) [338] were found encapsulated in the brain of patients with neurological symptoms. Such erratic parasitism may occur in immunocompromised patients [4]. In experimental mice, *M. yokogawai* worms were found to have invaded the submucosa of the small intestine when they were immunosuppressed [333].

Neodiplostomum seoulense

The normal habitat of this fluke is the duodenum in experimental rats and mice [230]. The major histopathological features of the affected mucosa include villous atrophy, crypt hyperplasia, mucosal inflammation, and bleeding [339]. The worms were found to entrap the host intestinal villi with their concave ventral curvature of the anterior body, piercing into the villous stroma with their tribocytic organ [339]. The affected villi undergo severe destruction with hemorrhages and inflammation, and finally the mucosa completely loses its integrity [339]. The histopathological changes due to *N. seoulense* are severer compared with those observed in other intestinal trematode infections including *Metagonimus*, *Pygidiopsis*, and *E. hortense* infections [4].

With respect to the pathogenicity of *N. seoulense*, the most striking feature is its high pathogenicity and even lethality to laboratory mice [340,341]. Almost all infected mice died by day 23 post-infection with 200 metacercariae per animal [341]. The infected mice revealed severely contracted intestines, strongly suggestive of an intestinal paralysis [341]. The fatality of the host animal varied according to genetic backgrounds of the mice [342]. Its tribocytic organ was suspected to be an important worm structure responsible for the host mucosal damage, because the tribocytic organ was shown to secrete alkaline phosphatase

tases, which could lyse the host villi and help the mucosal invasion of worms [343]. The tribocytic organ was also shown to contain neutral mucopolysaccharides, and thus the organ is suggested to play a protective role against host digestive enzymes [343].

IMMUNITY

Echinostomes

Immunological studies have been extensively performed in animal infecting echinostome species, i.e., *E. caproni*, *E. trivolvis*, and *Echinostoma friedi* [311]. Among the human infecting echinostomes, *E. hortense* has been mostly frequently studied [5,344,345]. Rodent hosts are known to express various types of resistance to echinostome infections [311]. One of them is spontaneous expulsion of primary infected worms [311]. However, the dynamics of worm expulsion are different depending on different parasite species and host species and strain [311, 344,346]. The elimination of *E. trivolvis* from mice occurs within 2-4 weeks [347], whereas worms can survive for long periods of time in golden hamsters [348]. In hamsters, *E. friedi* survive for at least 12 weeks, whereas the infection is expelled at 3-4 weeks in rats [349]. *E. caproni* worms are expelled from rats in 6-8 weeks, but they produce chronic infections in hamsters and mice [350]. An example of different immune responses by host strain includes *E. hortense* infection in BALB/c and C3H/HeN mice; whereas BALB/c mice reject worms, C3H/HeN mice undergo a chronic infection [120].

It has been suggested that 2 types of effector cells, i.e., goblet cells [346,351] and mucosal mast cells [352], are mainly associated with the worm expulsion [5]. However, the role of mast cells remains controversial [311]. For example, mastocytosis was suppressed in athymic nude mice but the kinetics of worm expulsion in these mice was similar to that of the conventional mice [353]. With respect to goblet cells, chronic infections with *E. caproni* in mice were accompanied by reduced goblet cell numbers [315,346,347]. In *E. trivolvis*-infected mice, the goblet cell numbers increased coinciding with worm expulsion, and expulsion was delayed when mice were treated with dexamethasone [353-355]. Similar immunosuppression effects by dexamethasone were observed in golden hamsters infected with *E. caproni* and *E. trivolvis* [356]. In addition, mucins increased by a primary *Nippostrongylus brasiliensis* infection in the intestine of mice were suggested to be responsible for a rapid expulsion of a challenge infection with *E. caproni* or *E. trivolvis* [357]. However,

a contradicting result was observed in the RAG-2-deficient mice, a strain of genetically lacking B- and T-lymphocytes, infected with *E. caproni*; goblet cell hyperplasia was marked in RAG-2-deficient mice but worms survived as they did in normal ICR mice [358].

Neutrophils and mononuclear inflammatory cells in the mesentery were suggested to be essential in determining the course of *E. caproni* infection in mice [359]. Some factors in mouse serum were also shown to induce retraction of collar spines in *E. trivolvis*, probably enhancing worm expulsion [360]. IgG, IgM, and IgA levels were elevated in serum and the small intestine of mice infected with *E. caproni* [361,362]. Antibodies in mouse serum covered the surface of *E. revolutum* in vitro, and worms obtained from immune mice were covered by antibodies [363]. Local immune responses seem to be more directly related with host resistance [5]. Whereas the serum level of IgG2a remained constant during the course of *E. caproni* infection, the intestinal IgG2a responses were related with worm expulsion [362]. Since IgG2a is an important element mediating T-helper-2 (Th2) immune responses of mice [364], the expulsion of echinostomes from the host intestine is suggested to be mediated by a Th2 immune response [362].

Th1 and Th2 cytokines profiles were studied in rodent hosts experimentally infected with *E. caproni* [365] or *E. hortense* [344, 345]. Mesenteric lymph node cells from *E. caproni*-infected mice, which are unable to expel worms in a short period of time, produced significantly higher levels of IFN- γ upon antigen stimulation in vitro during 3 wk after infection [365]. When these mice were injected with anti-IFN- γ monoclonal antibodies, the worm burden of *E. caproni* was significantly decreased [365]. It was thus suggested that in *E. caproni*-mouse model, Th1 responses mediated by IL-12, TNF- α , and IFN- γ are related to establishment of a chronic infection, whereas Th2 responses mediated by IL-4 and IL-5 lead to host protection and worm rejection [365,366]. Similarly, in *E. hortense* infection, BALB/c mice, a resistant mouse strain [344], exhibited stronger mRNA expressions of IL-4 and IL-5 in the spleen, than C3H/HeN mice, a susceptible strain which undergo a chronic infection [345]. Treatment of BALB/c mice with kitotifen, an anti-allergic drug, suppressed the mRNA levels of IL-4 and IL-5 and elevated the worm recoveries from this mouse strain [345].

Fasciolopsis buski

No information is available regarding immunity of the host against *F. buski* infection.

Gymnophalloides seoi

The susceptibility of mice to *G. seoi* infection is variable among different species, and even strains, of animals [164]. Among mouse strains, C3H/HeN mice were the most highly susceptible and the best for growth and development of the worms; although the worm recovery at days 7-21 post-infection was not sufficiently high [164,367]. However, when C3H/HeN mice were immunosuppressed by prednisolone injection, the survival and the recovery of the worms were greatly enhanced, and the enhancement was strongly correlated with the duration of immunosuppression [164,367]. Immunosuppression also enhanced the fecundity of the worms [164,367]. On the other hand, ICR and BALB/c mice are quite resistant against *G. seoi* infection [4]. ICR and BALB/c mice retain many worms by day 3 post-infection; however, most worms are expelled by day 7 post-infection [4,164]. It is of note that goblet cell proliferation was marked in the small intestines of infected mice, in particular, on the villous epithelia of the jejunum during days 3-7 post-infection [322]. The importance of goblet cells in worm expulsion was further suggested by only a small number of flukes retained in the intestines at days 7-14 post-infection [367,368]. This was supported by the minimal degree of goblet cell hyperplasia in immunocompromised mice, with many more flukes surviving in the intestine [367,368]. The goblet cell proliferation was shown to be highly dependent on CD4+ T-helper cells; however, both T-cell dependent and independent mechanisms seem to operate for expulsion of *G. seoi* worms from the mouse intestines [369].

Metagonimus and other heterophyids

Host protective mechanisms against *M. yokogawai* and other heterophyid flukes seem to be significant viewing from the fact that intestinal histopathology caused by these fluke infection is normalized around 3-4 wk after the infection [333]. One of the possible immune effectors includes intestinal intraepithelial lymphocytes (IEL) that increase remarkably along the villous epithelial layer of the infected rats and mainly consist of CD8+ cytotoxic T-cells [370]. Another effector cell is the lamina propria lymphocytes (LPL) that is chiefly consisted with IgA producing B-cells [370]. Mucosal mast cells were suggested as a third effector responsible for the worm expulsion from infected rats [371]. Goblet cells are a fourth effector for the expulsion of worms [2]. Immunogold studies have revealed that the antigenicity of *M. yokogawai* is originating from the syncytial tegument, tegumental cell cytoplasm, vitelline cells, and epithelial lamel-

lae of the cecum [372,373].

Neodiplostomum seoulense

Host protective immunity and survival of worms in the host intestine is known to be variable depending on different strains of mice [374]. BALB/c mice revealed a significantly higher recovery of worms than C3H mice, based on 28 days observation after an experimental infection [374]. Proliferation of mast cells was suggested to be a significant local immune response due to the presence of worms [375]. Goblet cells were also shown to increase significantly following experimental infection in mice [374]. However, these cells were not directly related with worm expulsion from the host intestine [374]. Binding of histamine released from mast cells to its receptors on intestinal smooth muscles would be more important than the level of histamine alone, or mastocytosis [376]. Recently, mixed Th1 and Th2 type immune responses were shown to occur during the course of *N. seoulense* infections in BALB/c mice, and IFN- γ (a Th1 cytokine), IL-4 (a Th2 cytokine), IgG, IgA, and IgG2a were all increased in the spleen and the small intestine [364]. Macrophages were also found significantly increased, which were found to kill *N. seoulense* worms in vitro [364].

Parasite-specific IgG levels were increased significantly in sera of infected mice, and 7 antigens, including 26, 30, 35, 43, 54, 67, and 94 kDa, were the major antigenic proteins of *N. seoulense* [377]. Serum and mucosal tissue IgA were increased after an experimental infection in mice, but the increase was not directly related to the worm expulsion [378]. Immunogold studies revealed that the tribocytic organ, seminal vesicle, caeca, and vitelline follicles were the major origins of worm antigens [379].

CLINICAL MANIFESTATIONS*Echinostomes*

Patients infected with echinostome species suffer from abdominal pain, diarrhea, tenesmus, easy fatigability, loss of body weight, and urinary incontinence [1,2,73,380]. The severity of symptoms is generally stronger than those seen in other intestinal fluke infections, such as heterophyids [5]. In this respect, reports of patients with *E. hortense* infection is noteworthy who suffered from severe epigastric discomfort and ulcerative lesions in the stomach or duodenum and diagnosed by recovery of worms at gastroduodenal endoscopy [125-129]. A patient admitted to a hospital because of epigastric pain and hematemesis revealed in gastroduodenoscopy an adult fluke attached at the

lesion, which was removed by an endoscopic clipper [125]. Another patient suffered from epigastric discomfort for a year, and acute epigastric pain and diarrhea for 1 month, which was accompanied by acidic belching, dizziness, and weight loss of 2 kg [126]. The third patient complained of epigastric discomfort for several days [127], and the fourth patient suffered from upper abdominal pain and discomfort persisting for 2 wk [128]. The fifth patient had epigastric discomfort, indigestion, anorexia, dizziness, headache, nausea, and vomiting; these symptoms became severer and she was hospitalized [129]. The clinical signs, including laboratory findings, in echinostomiasis are poorly known [5]. However, the levels of peripheral blood eosinophilia in *E. hortense* infection were dependent upon individual worm burdens; 11-24% (av. 17%) among the patients with more than 100 worms, 4-21% (av. 10%) among those with 51-100 worms, and 2-14% (av. 5%) among those with less than 50 worms [124]. Patients infected with *E. ilocanum* experienced intestinal colic and diarrhea [12]. Clinical symptoms in *E. cinetorchis*, *E. japonicus*, and *A. tyosenense* infections are not well known [5,7,56,92]. In 10 of 18 American tourists to Kenya and Tanzania who were echinostome egg positive in fecal examinations (adult flukes were not recovered), moderately severe abdominal cramps and loose or watery stools were encountered [381].

In gastroduodenal endoscopy, a moving, leaf-like *E. hortense* worm attached to an ulcerated mucosal layer of the distal part of the stomach was found [125]. The lesion was accompanied by a stage IIc or stage III early gastric cancer, and multiple ulcerations with bleeding were noted in the stomach and duodenum [125]. These ulcerations were suggested to be caused by the flukes [125]. Upper endoscopy of another patient revealed 2 *E. hortense* worms in the stomach; 1 worm was penetrating into the mucosa on the lesser curvature of the antrum; the other was attached to the mucosal surface of the greater curvature of the antrum [126]. The worm wiggled on stimulation and became tightly stretched during retrieval with a forcep [126]. Two more patients diagnosed at gastroduodenal endoscopy had 2 living worms each in the duodenal bulb area, at times accompanied by mucosal erosions [127,128]. In another patient, from whom 2 living *E. hortense* flukes were recovered, the affected gastric mucosa revealed signs of chronic gastritis, and at the proximal part of the duodenum multiple cured ulcerative lesions were observed [129]. In Japan, 4 cases of *E. hortense* infection were reported by recovery of flukes by upper endoscopy [382].

Fasciolopsis buski

Clinical manifestations remarkably vary from subclinical to fatal outcome [142]. In light infections, anemia, eosinophilia, headache, dizziness, gastric pain, and loose stools can occur [383]. In moderate and heavy infections, severe epigastric pain, abdominal pain, diarrhea, bowel obstruction, nausea, acute ileus, anasarca, and marked eosinophilia and leucocytosis may occur [383]. Due to allergic reactions to toxic worm metabolites, people infected with *F. buski* may suffer from facial, orbital edema, and anasarca. In heavy infections, intestinal perforation may occur [321].

Gymnophalloides seoi

Variable degrees of gastrointestinal troubles and indigestion, together with fever, anorexia, weight loss, easy fatigability, and weakness may occur in *G. seoi*-infected patients [149,150]. However, the degree of symptoms seems to be variable from individual patients [4]. A patient underwent acute pancreatitis and acute cholecystitis with episodes of epigastric discomforts, indigestion, and diarrhea [148]. Laboratory studies revealed elevated serum and urine amylase levels, increased serum alkaline phosphatase activity, and slight to moderate degrees of eosinophilia [148]. Five days after a treatment with praziquantel, epigastric pain and diarrhea completely disappeared, and serum and urine amylase levels returned to their normal levels [148]. Most other patients complained of only mild gastrointestinal troubles, such as indigestion [149,150]. However, it is noteworthy that 2 patients were accompanied by diabetes mellitus with high glucose levels [324]. Therefore, some relations between the *G. seoi* infection and diabetes were suspected [4]. In fact, some of the infected people living in a highly endemic area had symptoms, such as thirst, polydipsia, and polyuria [384], that may occur among diabetic patients.

Metagonimus and other heterophyids

Clinical symptoms due to *M. yokogawai* or other heterophyid infections are generally mild and transient unless heavily infected or immunocompromised [2,4]. In patients infected with *M. yokogawai*, the most frequent symptoms are mild to moderate degrees of abdominal pain, diarrhea, lethargy, anorexia, and weight loss [220,385]. The degree of clinical symptoms seems to be related to the individual worm burdens; heavier infection cases tend to suffer from more severe illness [4]. However, the severity of the symptoms may also be related to the susceptibility as well as acquired immunity of individual patient [4]. A vis-

itor to an endemic area, for example, suffered from a severe illness after a primary infection [386], whereas long-term residents generally complained of milder symptoms than expected [385]. Even a man living in a highly endemic area from whom as many as 63,587 worms were recovered complained of only minor gastrointestinal troubles, i.e., indigestion and epigastric pain [385]. Clinical symptoms due to other heterophyid fluke infections are not much different from those seen in *M. yokogawai* infection [4,151,152,387].

Neodiplostomum seoulense

In patients infected with *N. seoulense*, acute abdominal pain, diarrhea, lethargy, fever, and weight loss may occur [4]. However, clinical symptoms due to *N. seoulense* infection are not well documented except in the first patient, who experienced severe abdominal pain that led to admission to an emergency room of a university hospital [4,388]. The severity of symptoms may be dependent upon the individual worm burdens, as well as acquired immunity of each individual [4]. Patients repeatedly infected may complain of milder symptoms than primary infected patients, because most of the solidiers infected during survival trainings were asymptomatic [233,234]. However, because *N. seoulense* was reported to be highly pathogenic and even lethal to mice [341], infected humans may also experience severe clinical course as seen in the first patient [388].

DIAGNOSIS

Echinostomes

Human echinostomiasis is diagnosed by recovery of eggs in fecal examinations [5]. Careful microscopic observations with measurements of the eggs can provide even specific diagnosis in known endemic areas of certain echinostome species [5]. However, recovery and identification of the adult flukes is strongly recommended, if a definite diagnosis is preferred [4,5]. The detectability of eggs in fecal examinations varies remarkably depending on different echinostome species [5]. The fecundity is considerably high in *E. hortense* [119] and *E. cinetorchis* [101], whereas it is very low in *E. japonicus* [60] and *A. tyosenense* [7]. The difference may be due to remarkably different numbers of intrauterine eggs and the different egg laying capacity of each echinostome species [4,5].

Fasciolopsis buski

Diagnosis is carried out by coprological examinations to detect

eggs or occasionally by examination of expelled adult worms vomited or passed in the stool [142]. Serodiagnosis detecting serum antibodies may help the diagnosis [389].

Gymnophalloides seoi

The diagnosis can be made by detection of eggs in the feces; however, an expert is needed to identify the eggs [4,158]. The eggs are very small, only 0.020-0.025 mm in length, smaller than those of *Clonorchis sinensis*, *Metagonimus yokogawai*, or other heterophyids, and have a very thin and transparent shell [148-150]. The size of *G. seoi* eggs is similar to that of *P. summa* [2]. The problem is that the eggs may be overlooked or misdiagnosed as an air bubble or other artifacts in routine fecal examinations performed by formalin-ether sedimentation or cellophane thick smear techniques [149]. Another problem is a very low egg laying capacity of *G. seoi*, i.e., only 2-84 eggs per adult fluke in the human host [384], compared to other intestinal parasites [2]. It is assumed that unless more than 100 worms are present, less than 8,400 eggs would be discharged in a whole-day stool, and the eggs per gram of feces (EPG) would be only 42 (daily stool weight; 200 g) [4]. This value means the appearance of only 1-2 eggs on the whole field of a fecal smear made by the Kato-Katz technique (41.7 mg of feces/smear) [2,4].

Metagonimus and other heterophyids

When heterophyid eggs are detected in fecal examinations, it is expressed using the term 'heterophyid fluke infection', because the eggs of different heterophyid species closely resemble each other [4,390]. The diagnosis can be confirmed after the recovery of adult flukes following anthelmintic treatment and purgation [5]. The specific diagnosis only by eggs is difficult in areas of no known endemicity, as well as in areas with mixed heterophyid infections [4].

However, close observations and measurements of heterophyid eggs are useful for differential diagnosis [4,390]. For example, *M. yokogawai* eggs can be differentiated from other heterophyid eggs by their length of 26.9-31.6 μ m, elliptical shape with length/width ratio of 1.5-2.1, clean shell surface, less prominent operculum, no shoulder rims, and dark yellow or brown color [390]. *M. takahashii* and *M. miyatai* eggs are similar to those of *M. yokogawai*, but larger; hence, the measurement of egg size is essential [4]. *H. nocens* eggs are similar to those of *M. yokogawai*, but the former is a little smaller and has slight attenuations at one or both ends [390]. *H. continua* eggs are broadly oval in shape, and difficult to differentiate from those of *M. yokogawai*

[390]. *P. summa* eggs are characteristically small and pyriform in shape [390]; they resemble the eggs of *Clonorchis sinensis*, a liver fluke, but lack muskmelon patterns on the shell surface, and shoulder rims around the operculum are inconspicuous [4]. Serological tests such as ELISA are helpful in false egg negative cases infected with *M. yokogawai* [386,391].

Eggs of *H. taichui*, *H. pumilio*, and *H. yokogawai* closely resemble each other, and need differentiation also from eggs of *Opisthorchis viverrini* and lecithodendriid flukes, i.e., *Phaneropsolus bonnei* and *Prosthodendrium molencampi* [4,243].

Neodiplostomum seoulense

The diagnosis of *N. seoulense* infection can be done by recovery of typical eggs in the feces [230]. The eggs are ellipsoid to elliptical, thin-shelled, with an inconspicuous operculum, and frequently asymmetrical [230]. They differ from the eggs of *E. hortense* or *E. cinetorchis*, in that they have a clean shell surface and, unlike the latter, they do not have abopercular wrinkles at the posterior end [4,5].

TREATMENT AND CONTROL

Echinostomes

Treatment of echinostome infections can be done using 10-20 mg/kg praziquantel in a single oral dose [5]. Albendazole may be effective [4]. For prevention and control, eating raw or improperly cooked fresh water fish and fresh or brackish water snails should be avoided [4,5].

Fasciolopsis buski

Tetrachlorethylene had been the drug of choice before introduction of praziquantel for treatment of fasciolopsiasis in 1983 [142]. Now praziquantel, at the dose of 15 mg/kg single dose, is prescribed even in severe fasciolopsiasis patients [142]. A new drug, triclabendazole, has been introduced showing a promising efficacy without side effects in pigs [142,392]. Two other new drugs, oxiclozanide and rafozanide, were nearly equally effective as triclabendazole in pigs [392].

Gymnophalloides seoi

For treatment of *G. seoi* infection in humans, praziquantel in a single oral dose of 10 mg/kg is prescribed [4,148,149]. Albendazole may also be effective against *G. seoi* infection, but this needs confirmation [4]. The best way to prevent *G. seoi* infection is the avoidance of consuming infected oysters, under raw

or improperly cooked conditions [4]. As control measures, oyster irradiation [393] and repeated chemotherapy of the people in endemic areas [384] were tried with considerable success.

Metagonimus and other heterophyids

For all infections with heterophyid flukes, praziquantel is the drug of choice [4]. A single oral dose of 10-20 mg/kg praziquantel is satisfactory, with 95-100% cure rate for *M. yokogawai* infection [394,395]. Niclosamide was used in vitro to observe damage to *H. taichui* worms [396]. Irradiation of the sweetfish, *Plecoglossus altivelis*, by 200 Gy was highly effective in controlling infectivity of the *M. yokogawai* metacercariae [397]. The heterophyid fluke infections could be prevented by the avoidance of eating uncooked fresh water or brackish water fish [4].

Neodiplostomum seoulense

Praziquantel in a single oral dose of 10-20 mg/kg is a highly effective treatment for *N. seoulense* infection [4,230]. For prevention, ingestion of raw or improperly cooked flesh of snakes or frogs should be avoided [4].

REFERENCES

1. Fried B, Graczyk TK, Tamang L. Food-borne intestinal trematodiasis in humans. *Parasitol Res* 2004; 93: 159-170.
2. Chai JY, Lee SH. Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* 2002; 51: 129-154.
3. Chai JY, Murrell KD, Lymbery AJ. Fish-borne parasitic zoonoses: status and issues. *Int J Parasitol* 2005; 35: 1233-1254.
4. Chai JY. Intestinal flukes. In Murrell KD, Fried B eds, *Food-borne Parasitic Zoonoses: Fish and Plant-borne Parasites*. World Class Parasites, Vol. 11. New York, USA. Springer. 2007, p 53-115.
5. Chai JY. Echinostomes in humans. In Fried B, Toledo R eds, *The Biology of Echinostomes*. New York, USA. Springer. 2009, p 147-183.
6. Yu SH, Mott KE. Epidemiology and morbidity of food-borne intestinal trematode infections. *Trop Dis Bull* 1994; 91: R125-R152.
7. Chai JY, Han ET, Park YK, Guk SM, Lee SH. *Acanthoparyphium tyosenense*: the discovery of human infection and identification of its source. *J Parasitol* 2001; 87: 794-800.
8. Yamaguti S. Studies on the helminth fauna of Japan. Part 25. Trematodes of birds, IV. *Jpn J Zool* 1939; 8: 131-210.
9. Kim YK, Yu JE, Chung EY, Chung PR. *Acanthoparyphium tyosenense* (Digenea: Echinostomatidae): experimental confirmation of the cercaria and its complete life history in Korea. *J Parasitol* 2004; 90: 97-102.
10. Ito J. Studies on the brackish water cercariae in Japan. III. Three new echinostome cercariae in Tokyo Bay, with a list of Japanese

- echinostome cercariae (Trematoda). *Jpn J Med Sci Biol* 1957; 10: 439-453.
11. Han ET, Kim JL, Chai JY. Recovery, growth and development of *Acanthoparyphium tyosenense* (Digenea: Echinostomatidae) in experimental chicks. *J Parasitol* 2003; 89: 176-179.
 12. Beaver PC, Jung RC, Cupp EW. *Clinical Parasitology*. 9th ed. Philadelphia, USA. Lea & Febiger. 1984.
 13. Maji AK, Bera DK, Manna B, Nandy A, Addy M, Bandyopadhyay AK. First record of human infection with *Echinostoma malayanum* in India. *Trans R Soc Trop Med Hyg* 1993; 87: 673.
 14. Radomyos B, Wongsaraj T, Wilraitana P, Radomyos P, Praewanich R, Meesomboon V, Jongsuksuntikul, P. Opisthorchiasis and intestinal fluke infections in northern Thailand. *Southeast Asian J Trop Med Public Health* 1998; 29: 123-127.
 15. Belizario VY, Geronilla GG, Anastacio MB, de Leon WU, Subaan AU, Sebastian AC, Bangs MJ. *Echinostoma malayanum* infection, the Philippines. *Emerg Infect Dis* 2007; 13: 1130-1131.
 16. Premvati G, Pande V. On *Artyfechinostomum malayanum* (Leiper, 1911) Mendheim, 1943 (Trematoda: Echinostomatidae) with synonymy of allied species and genera. *Proceed Helminthol Soc Washington* 1974; 41: 151-160.
 17. Kostadinova A, Vaucher C, Gibson DI. Redescriptions of two echinostomes from birds in Paraguay, with comments on *Drepanocephalus* Diet, 1909 and *Paryphostomum* Diez, 1909 (Digenea: Echinostomatidae). *Syst Parasitol* 2002; 53: 147-158.
 18. Lane C. *Artyfechinostomum sufrartyfex*. A new parasitic echinostome of man. *Indian J Med Res* 1915; 2: 977-983.
 19. Reddy DG, Varmah K. *Paryphostomum sufrartyfex* (intestinal fluke) infection in man. *Ind Med Gaz* 1950; 85: 546-547.
 20. Ahluwalia SS. Studies on some helminths of the domestic pigs (*Sus scrofa domestica*) in western Uttar Pradesh. *J Helminthol* 1962; 36: 347-364.
 21. Dubey JP, Srivastava HO, Sahasrabudhe VK. *Artyfechinostomum sufrartyfex* infection in cat and dog in India. *Trop Geogr Med* 1969; 21: 210-213.
 22. Raghunathan VS, Srinivasan T. *Artyfechinostomum mehrai* infestation. A case report. *J Indian Med Assoc* 1962; 38: 485-487.
 23. Lie KJ. Studies on Echinostomatidae in Malaya. III. The adult *Echinostoma malayanum* Leiper, 1911 (Trematoda) and the probable synonymy of *Artyfechinostomum sufrartyfex* Lane, 1915. *Z Parazitenk* 1963; 23: 124-135.
 24. Bandyopadhyay AK, Nandy A. A preliminary observation on the prevalence of echinostomes in a tribal community near Calcutta. *Ann Trop Med Parasitol* 1986; 80: 373-375.
 25. Bandyopadhyay AK, Manna B, Nandy A. Human infection of *Artyfechinostomum oraoni* n. sp. (Paryphostominae: Echinostomatidae) in a tribal community, "Oraons" in West Bengal, India. *Indian J Parasitol* 1989; 13: 191-196.
 26. Maji AK, Manna B, Bandyopadhyay AK, Bera DK, Addy M, Nandy A. Studies on the life cycle of *Artyfechinostomum oraoni* Bandyopadhyay, Manna & Nandy, 1989: embryogenesis and development in the intermediate host. *Indian J Med Res* 1995; 102: 124-128.
 27. Bandyopadhyay AK, Maji AK, Manna B, Bera DK, Addy M, Nandy A. Pathogenicity of *Artyfechinostomum oraoni* in naturally infected pigs. *Trop Med Parasitol* 1995; 46: 138-139.
 28. Jueco NL, Monzon RB. *Cathaemacia cabrerai* sp. n. (Trematoda: Cathaemaciidae) a new oarqasite of man in the Philippines. *Southeast Asian J Trop Med Public Health* 1984; 15: 427-429.
 29. Tanabe H. Studien uber die Trematoden mit Susswasser-fischen als Zwischenwirt. I. *Stamnosoma armatum* n. g., n. sp. *Kyoto Igaku Zasshi* 1922; 19: 239-252.
 30. Hong SJ, Seo BS, Lee SH, Chai JY. A human case of *Centrocestus armatus* infection in Korea. *Korean J Parasitol* 1988; 26: 55-60.
 31. Takahashi S. A contribution to the life-history of *Stamnosoma armatum* Tanabe. *Okayama Igakkai Zasshi* 1929; 4: 1759-1771.
 32. Lee JK, Lee HI, Baek BK, Kim PG. Survey on encysted cercariae of trematodes from fresh-water fishes in Tongjin riverside areas in Korea. *Korean J Parasitol* 1984; 22: 190-202.
 33. Hong SJ, Woo HC, Kim IT. Study on *Centrocestus armatus* in Korea I. Infection status of *Zacco platypus* and *Z. temminckii* with the metacercariae of *C. armatus*. *Korean J Parasitol* 1989; 27: 41-46.
 34. Ryang YS, Ahn YK, Yoon MB. Trematode infections in the small intestine of *Egretta alba medesta* in Kanhwon-do. *Korean J Parasitol* 1991; 29: 227-233.
 35. Sohn WM, Chai JY. Infection status with helminthes in feral cats purchased from a market in Busan, Republic of Korea. *Korean J Parasitol* 2005; 43: 93-100.
 36. Han ET, Shin EH, Phommakorn S, Sengvilaykham B, Kim JL, Rim HJ, Chai JY. *Centrocestus formosanus* (Digenea: Heterophyidae) encysted in the freshwater fish, *Puntius brevis*, from Lao PDR. *Korean J Parasitol* 2008; 46: 49-53.
 37. Waikagul J, Wongsaraj T, Radomyos P, Meesomboon V, Praewanich R, Jongsuksuntikul P. Human infection of *Centrocestus caninus* in Thailand. *Southeast Asian J Trop Med Public Health* 1997; 28: 831-835.
 38. Saenphet S, Wongsawad C, Saenphet K, Rojanapaibul A, Vanitanakom P, Chai JY. Chronological observations of intestinal histopathology in rats (*Rattus norvegicus*) infected with *Centrocestus caninus*. *Southeast Asian J Trop Med Public Health* 2006; 37(suppl 3): 69-73.
 39. Nishigori M. On a new trematode, *Stamnosoma formosanum* n. sp., and its life history. *Taiwan Igakkai Zasshi* 1924; 234: 181-228.
 40. Ito J. *Metagonimus* and other human heterophyid trematodes. *Progress of Medical Parasitology in Japan* 1964; 1: 314-393.
 41. Chen HT. The metacercaria and adult of *Centrocestus formosanus* (Nishigori, 1924), with notes on the natural infection of rats and cats with *C. armatus* (Tanabe, 1922). *J Parasitol* 1942; 28: 285-298.
 42. Martin WE. The life cycle of some Hawaiian heterophyid trematodes. *J Parasitol* 1958; 44: 305-321.
 43. Premvati G, Pande V. On *Centrocestus formosanus* (Nishigori, 1924) Price, 1932 and its experimental infections in white leghorn chicks. *Jpn J Parasitol* 1974; 23: 79-84.

44. Madhavi R. Distribution of metacercariae of *Centrocestus formosanus* (Trematoda: Heterophyidae) on the gills of *Aplocheilichthys* (Trematoda: Heterophyidae) on the gills of *Aplocheilichthys pan-chax*. *J Fish Biol* 1986; 29: 685-690.
45. Scholz T, Salgado-Maldonado G. The introduction and dispersal of *Centrocestus formosanus* (Nishigori, 1924) (Digenea: Heterophyidae) in Mexico: a review. *American Midland Naturalist* 1999; 143: 185-200.
46. Rim HJ, Sohn WM, Yong TS, Eom KS, Chai JY, Min DY, Lee SH, Hoang EH, Phommasack B, Insisengmay S. Fishborne trematode metacercariae detected in freshwater fish from Vientiane Municipality and Savannakhet Province, Lao PDR. *Korean J Parasitol* 2008; 46: 253-260.
47. Kurokawa T. On a new species of *Stamnosoma* found from a man. *Tokyo Iji Shinshi* 1935; 2915: 293-298.
48. Yamaguti S. Part I. The digenetic trematodes of vertebrates. In *Systema Helminthum* 1958; Vol. I: 1-979. New York, USA. Interscience Publishers Inc.
49. Chen CE, Cai GD. Morphological testimony: demonstration of a case of human infestation with *Cotylurus japonicus* Ishii, 1932 (Trematoda, Strigeidae). *Bull Hunan Med Coll* 1985; 10: 31-34.
50. Cheng YZ, Lin JX, Chen BJ, Fang YY, Guo ZF, Xu GF, Wu ZY. A new species of *Echinochasmus* parasitic in man and observation of its experimental infection. *Wuyi Sci J* 1992; 9: 43-48.
51. Cheng YZ, Zhang Y, Lin C, Zhan Z, Yu X, Lin JX, Cai MJ, Che W. Taxonomic studies on *Echinochasmus fujianensis* and its related species by random amplified polymorphic DNA analysis and experimental infection. *Chinese J Parasitol Parasit Dis* 1999; 17: 135-139.
52. Tanabe H. Studies on trematodes with fresh water fishes as their intermediate host III. On a new species, *Echinochasmus japonicus* (n.sp.). *Nippon Byori Gakkai Kaishi* 1926; 16: 295-296.
53. Ujiiie N. On the development, the structure of *Echinochasmus japonicus* and its parasitism in man. *Taiwan Igakkai Zasshi* 1939; 35: 535-546.
54. Lin JX, Chen YZ, Liang CZ, Yang LB, Zhuang HJ. Epidemiological investigation and experimental infection with *Echinochasmus japonicus*. *Chinese J Parasitol Parasit Dis* 1985; 3: 89-91.
55. Zhu D, Lin J, Cheng Y, Liang C, Wang X, Guo Z, Zhuang H. Pyquion treatment of *Echinochasmus japonicus* infection. *Chinese J Parasitol Parasit Dis* 1986; 4: 1-4.
56. Seo BS, Lee SH, Chai JY, Hong SJ. Studies on intestinal trematodes in Korea XX. Four cases of natural human infection by *Echinochasmus japonicus*. *Korean J Parasitol* 1985; 23: 214-220.
57. Lee JK, Lee HI, Baek BK, Kim PG. Survey on encysted cercariae of trematodes from fresh-water fishes in Mangyeong riverside area. *Korean J Parasitol* 1983; 21: 187-192.
58. Choi MH, Kim SH, Chung JH, Jang HJ, Eom JH, Chung BS, Sohn WM, Chai JY, Hong ST. Morphological observations of *Echinochasmus japonicus* cercariae and the in vitro maintenance of its life cycle from cercariae to adults. *J Parasitol* 2006; 92: 236-241.
59. Lee JK, Lee HI, Baek BK, Kim PG. Survey on encysted cercariae of trematodes from fresh-water fishes in Tongjin riverside areas in Korea. *Korea J Parasitol* 1984; 22: 190-202.
60. Chai JY, Hong SJ, Son DW, Lee SH, Seo BS. Metacercariae of the *Echinochasmus japonicus* encysted in a fresh water fish, *Pseudorasbora parva*, and their development in experimental mice. *Korean J Parasitol* 1985; 23: 221-229.
61. Eom KS, Rim HJ. A study on the parasitic helminths of domestic duck (*Anas platyrhynchos* var. *domestica* Linnaeus) in Korea. *Korean J Parasitol* 1984; 22: 215-221.
62. Chai JY, Park JH, Jung BK, Guk SM, Kim JL, Shin EH, Klein TA, Kim HC, Chong ST, Baek LJ, Song JW. Echinostome infections in the striped-field mouse, *Apodemus agrarius*, and the Ussuri white-toothed shrew, *Crocidura lasiura*, caught near the demilitarized zone, Gyonggi-do (province), republic of Korea. *Korean J Parasitol* 2009; 47: 311-314.
63. Liang C, Ke XL. *Echinochasmus jiufoensis* sp. nov., a human parasite from Guangzhou (Trematoda: Echinostomatidae). *Acta Zootaxonomica Sinica* 1988; 13: 4-8.
64. Witenberg G. On two parasite species of the genus *Echinochasmus* (Trematoda) in dogs and cats in Palestine. *Z Parazitenk* 1932; 5: 3213-3216.
65. Xiao X, Wang TP, Lu DB, Gao JF, Xu LF, Wu WD, Mei JD, Xu FN. The first record of human infection with *Echinochasmus liliputanus*. *Chinese J Parasitol Parasit Dis* 1992; 10: 132-135.
66. Xiao X, Wang T, Zheng X, Shen G, Tian Z. In vivo and in vitro encystment of *Echinochasmus liliputanus* cercariae and biological activity of the metacercariae. *J Parasitol* 2005; 91: 492-498.
67. El-Gayar AK. Studies on some trematode parasites of stray dogs in Egypt with a key to the identification of intestinal trematodes of dogs. *Vet Parasitol* 2007; 144: 360-365.
68. Xiao X, Dabing L, Tianping W, Jianfeng G, Chuang Z, Banju Z, Jingshen A, Hongchuan P, Mingrong X, Weido W, Jide M, Xu F. Studies on mode of human infection with *Echinochasmus liliputanus*. *Chinese J Parasitol Parasit Dis* 1995; 13: 197-199.
69. Tanabe H. Studies on trematodes with freshwater fish as its intermediate host. II. On *Echinochasmus perfoliatus* (Ratz) found in Japan. *Okayama Igakkai Zasshi* 1922; No. 387: 175-195.
70. Shimalov VV, Shimalov VT. Human parasitizing echinostomatata (Trematoda, fasciolidae) in mammals in Belorussia forest zone. *Meditsinskaia Parazitologiya i Parazitamy Bolezni*, No. 1 2002; 47-50.
71. Saeed I, Maddox-Hyttel C, Monrad J, Kapel CM. Helminths of red foxes (*Vulpes vulpes*) in Denmark. *Vet Parasitol* 2006; 139: 168-179.
72. Hirazawa I. On *Echinochasmus perfoliatus* (Ratz) found in man. *Tokyo Iji Shinshi* 1928; 2577: 1328-1334.
73. Rim HJ. Echinostomiasis. *CRC Handbook Series in Zoonoses, Section C: Parasitic Zoonoses, Vol. III. Trematode Zoonoses*. Boca Raton, Florida, USA. CRC Press Inc. 1982, p 53-69.
74. Yamashita J. Echinostome. *Progress of Med Parasitol in Japan* 1964; Vol. 1: 287-313. Tokyo, Japan. Meguro Parasitological Museum.
75. Lee SH, Sohn WM, Chai JY. *Echinostoma revolutum* and *Echinoparyphium recurvatum* recovered from house rats in Yangyang-

- gun, Kangwon-do. *Korean J Parasitol* 1990; 28: 235-240.
76. Betlejewska KM, Jorol EN. Taxonomic, topical and quantitative structure of the community of intestinal flukes (Digenea) of mallards, *Anas platyrhynchos* Linnaeus, 1758 from the area of Szczecin. *Wiadomosci Parazytologiczne* 2002; 48: 343-357.
 77. Ried B. Biology of echinostomes except *Echinostoma*. *Adv Parasitol* 2001; 49: 163-210.
 78. Evans NA, Gordon DM. Experimental observations on the specificity of *Echinoparyphium recurvatum* toward second intermediate hosts. *Z Parazitenk* 1983; 69: 217-222.
 79. McCarthy AM. The influence of second intermediate host dispersion pattern upon the transmission of cercariae of *Echinoparyphium recurvatum* (Digenea: Echinostomatidae). *Parasitology* 1990; 101: Pt 1: 43-47.
 80. Sohn WM. Life history of *Echinoparyphium recurvatum* (Trematoda: Echinostomatidae) in Korea. *Korean J Parasitol* 1998; 36: 91-98.
 81. Yurlova NI, Vodyanitskaya SN, Serbina EA, Biserkov VY, Georgiev BB, Chipev NH. Temporal variation in prevalence and abundance of metacercariae in the pulmonate snail *Lymnaea stagnalis* in Chany Lake, West Siberia, Russia: Long-term patterns and environmental covariates. *J Parasitol* 2006; 92: 249-259.
 82. Fain A, Galal A. Identification of the trematode *Echinoparyphium recurvatum* (von Linstow, 1873) in man and in a rodent from Egypt. *Ann Soc Belg Med Trop* 1977; 57: 583-587.
 83. Cheng YZ, Lin JX, Fang YY, Lin AQ, Chen BJ, Zhuang HJ. Discovery of human infection with *Echinostoma angustitestis*. *Chinese Journal of Zoonoses* 1992; 8: 7-8.
 84. Ando R, Ozaki Y. On four new species of trematodes of the family Echinostomatidae. *Dobutsugaku Zasshi* 1923; 35: 108-119.
 85. Sugimoto M. On a trematode parasite (*Echinostoma cinetorchis* Ando et Ozaki, 1923) from a Formosan dog. *Nippon Juigaku Zasshi* 1933; 12: 231-236.
 86. Fischthal JH, Kuntz RE. Some digenetic trematodes of mammals from Taiwan. *Proceed Helminthol Soc Washington* 1975; 42: 149-157.
 87. Seo BS, Rim HJ, Lee CW. Studies on the parasitic helminths of Korea I. Trematodes of rodents. *Korean J Parasitol* 1964; 2: 20-26.
 88. Seo BS, Cho SY, Hong ST, Hong SJ, Lee SH. Studies on parasitic helminths of Korea V. Survey on intestinal trematodes of house rats. *Korean J Parasitol* 1981; 19: 131-136.
 89. Wu G. Human parasitology. Beijing, China. Rin Min Hu Sheng, 2004.
 90. Takahashi S, Ishii T, Ueno N. A human case of *Echinostoma cinetorchis*. *Tokyo Iji Shinshi* 1930; 2757: 141-144.
 91. Kawahara S, Yamamoto E. Human cases of *Echinostoma cinetorchis*. *Tokyo Iji Shinshi* 1933; No. 2840: 1794-1796.
 92. Seo BS, Cho SY, Chai JY. Studies on intestinal trematodes in Korea I. A human case of *Echinostoma cinetorchis* infection with an epidemiological investigation. *Seoul J Med* 1980; 21: 21-29.
 93. Ryang YS, Ahn YK, Kim WT, Shin KC, Lee KW, Kim TS. Two cases of human infection by *Echinostoma cinetorchis*. *Korean J Parasitol* 1986; 24: 71-76.
 94. Lee SK, Chung NS, Ko IH, Ko HI, Sohn WM. A case of natural human infection by *Echinostoma cinetorchis*. *Korean J Parasitol* 1988; 26: 61-64.
 95. Lee SH, Chai JY, Hong ST, Sohn WM. Experimental life history of *Echinostoma cinetorchis*. *Korean J Parasitol* 1990; 28: 39-44.
 96. Chung PR, Jung Y, Park YK. Segmentina hemisphaerula: a new molluscan intermediate host for *Echinostoma cinetorchis* in Korea. *J Parasitol* 2001; 87: 1169-1171.
 97. Ahn YK, Ryang YS, Chai JY, Sohn WM. Cercarial shedding of *Echinostoma cinetorchis* and experimental infection of the cercariae to several kinds of snails. *Korean J Parasitol* 1989; 27: 203-211.
 98. Chung PR, Jung Y. *Cipangopaludina chinensis malleata* (Gastropoda: Viviparidae): a new second molluscan intermediate host of a human intestinal fluke *Echinostoma cinetorchis* (Trematoda: Echinostomatidae) in Korea. *J Parasitol* 1999; 85: 963-964.
 99. Ando R, Tsuyuki H. Studies on intestinal parasites taking rats as their final hosts. II. On the trematode *Echinostoma* and its second intermediate host in Nagoya city. *Tokyo Iji Shinshi* 1923; 2340: 1487-1499.
 100. Saito S, Tani S. Comparison of metacercariae of *Echinostoma hortense* Asada, 1926 and *Echinostoma cinetorchis* Ando et Ozaki, 1923 in loach, *Misgurnus anguillicaudatus*. *Jpn J Parasitol* 1982; 31: 281-287.
 101. Seo BS, Park YH, Chai JY, Hong SJ, Lee SH. Studies on intestinal trematodes in Korea. XIV. Infection status of loaches with metacercariae of *Echinostoma cinetorchis* and their development in albino rats. *Korean J Parasitol* 1984; 22: 181-189.
 102. Cho SY, Kang SY, Ryang YS. Helminthes infections in the small intestine of stray dogs in Eunjungbu City, Kyunggi Do, Korea. *Korean J Parasitol* 1981; 19: 55-59.
 103. Lee SH, Lee JK, Sohn WM, Hong ST, Hong SJ, Chai JY. Metacercariae of *Echinostoma cinetorchis* encysted in the fresh water snail, *Hippeutis (Helicorbis) cantori*, and their development in rats and mice. *Korean J Parasitol* 1988; 26: 189-197.
 104. Kanev I. Life-cycle, delimitation and redescription of *Echinostoma revolutum* (Froelich, 1982) (Trematoda: Echinostomatidae). *Syst Parasitol* 1994; 28: 125-144.
 105. Huffman JE, Fried B. *Echinostoma* and echinostomiasis. *Adv Parasitol* 1990; 29: 215-269.
 106. Kostadinova A. *Echinostoma echinatum* (Zeder 1803) sensu Kanev (Digenea: Echinostomatidae): a note of caution. *Syst Parasitol* 1995; 32: 23-26.
 107. Kostadinova A, Gibson DI. The systematic of the echinostomes. In Fried B, Graczyk TK eds, *Echinostomes as Experimental Models for Biological Research*. Dordrecht, Boston, London. Kluwer Academic Publishers 2000, p 31-57.
 108. Fried B, Graczyk TK. Recent advances in the biology of *Echinostoma* species in the 'revolutum' group. *Adv Parasitol* 2004; 58: 139-195.
 109. Sandground JH, Bonne C. *Echinostoma lindoensis* n. sp., a new parasite of man in the Celebes with an account of its life histo-

- ry and epidemiology. *Am J Trop Med* 1940; 20: 511-536.
110. Lie KJ. Further studies on the life history of *Echinostoma lindoense* Sandground and Bonne, 1940 (Trematoda: Echinostomatidae) with a report of its occurrence in Brazil. *Proceedings of the Helminthol Soc Washington* 1968; 35: 74-77.
 111. Asada J. On a new echinostomatid trematode and its life history. *Trans Jpn Soc Pathol* 1926; 16: 293-294.
 112. Park JT. A rat trematode, *Echinostoma hortense* Asada, from Korea. *Keijo J Med* 1938; 9: 283-286.
 113. Fan SQ, Sun MF. *Echinostoma hortense*. *Bull Harbin Med Univ* 1989; 23: 161.
 114. Kostadinova A, Gibson DI. Isthmiophora Lühe, 1909 and *Euparyphium* Diet, 1909 (Digenea: Echinostomatidae) re-defined, with comments on their nominal species. *Syst Parasitol* 2002; 52: 205-217.
 115. Ahn YK, Ryang YS. Epidemiological studies on *Metagonimus* infection along the Hongcheon river, Kangwon province. *Korean J Parasitol* 1988; 26: 207-213.
 116. Chai JY, Hong SJ, Sohn WM, Lee SH, Seo BS. Studies on intestinal trematode in Korea. XVI. Infection status of loaches with the metacercariae of *Echinostoma hortense*. *Korean J Parasitol* 1985; 23: 18-23.
 117. Ryang YS, Ahn YK, Lee KW, Kim TS, Hhan MH. Two cases of natural human infection by *Echinostoma hortense* and its second intermediate host in Wonju area. *Korean J Parasitol* 1985; 23: 33-40.
 118. Ryang YS. Studies on *Echinostoma* spp. in the Chungju Reservoir and upper streams of the Namhan River. *Korean J Parasitol* 1990; 28: 221-233.
 119. Seo BS, Chun KS, Chai JY, Hong SJ, Lee SH. Studies on intestinal trematodes in Korea XVII. Development and egg laying capacity of *Echinostoma hortense* in albino rats and human experimental infection. *Korean J Parasitol* 1985; 23: 24-32.
 120. Lee KJ, Park SK, Im JA, Kim SK, Kim GH, Kim GY, Yang EJ, Ryang YS. Susceptibility of several strains of mice to *Echinostoma hortense* infection. *Korean J Parasitol* 2004; 42: 51-56.
 121. Seo BS, Hong ST, Chai JY, Lee SH. Studies on intestinal trematodes in Korea VII. A human case of *Echinostoma hortense* infection. *Korean J Parasitol* 1983; 21: 219-223.
 122. Miyamoto K, Nakao M, Inaoka T. Studies on the zoonoses in Hokkaido, Japan. 5. On the epidemiological survey of *Echinostoma hortense* Asada, 1926. *Jpn J Parasitol* 1983; 32: 261-269.
 123. Chen XQ, Feng GX, Qian ZF. Survey on infection due to *Echinostoma hortense* in Liaoning Province. *Chinese J Parasitol Parasit Dis* 1993; 11: 226.
 124. Lee SK, Chung NS, Ko IH, Sohn WM, Hong ST, Chai JY, Lee SH. An epidemiological survey of *Echinostoma hortense* infection in Chongsong-gun, Kyongbuk Province. *Korean J Parasitol* 1988; 26: 199-206.
 125. Chai JY, Hong ST, Lee SH, Lee GC, Min YI. A case of echinostomiasis with ulcerative lesions in the duodenum. *Korean J Parasitol* 1994; 32: 201-204.
 126. Lee OJ, Hong SJ. Gastric echinostomiasis diagnosed by endoscopy. *Gastrointest Endosc* 2002; 55: 440-442.
 127. Cho CM, Tak WY, Kweon YO, Kim SK, Choi YH, Kong HH, Chung DI. A human case of *Echinostoma hortense* (Trematoda: Echinostomatidae) infection diagnosed by gastroduodenal endoscopy in Korea. *Korean J Parasitol* 2003; 41: 117-120.
 128. Chang YD, Sohn WM, Ryu JH, Kang SY, Hong SJ. A human infection of *Echinostoma hortense* in duodenal bulb diagnosed by endoscopy. *Korean J Parasitol* 2005; 43: 57-60.
 129. Park CJ, Kim J. A human case of *Echinostoma hortense* infection diagnosed by endoscopy in area of southwestern Korea. *Korean J Med* 2006; 71: 229-234.
 130. Cross JH, Bassaca-Sevilla V. Studies on *Echinostoma ilocanum* in the Philippines. *Southeast Asian J Trop Med Public Health* 1986; 17: 23-26.
 131. Radomyos P, Bunnag D, Harinasuta T. *Echinostoma ilocanum* (Garrison, 1908) Odhner, 1911, infection in man in Thailand. *Southeast Asian J Trop Med Public Health* 1982; 13: 265-269.
 132. Cross JH, Bassaca-Sevilla V. Intestinal parasitic infections in Southeast Asia. *Southeast Asian J Trop Med Public Health* 1981; 12: 262-274.
 133. Grover M, Dutta R, Kumar R, Aneja S, Mehta G. *Echinostoma ilocanum* infection. *Indian Pediatr* 1998; 35: 549-552.
 134. Majima M. On *Echinostoma macrorchis* found in man. *Kumamoto Igakkai Zasshi* 1927; 2252: 2260.
 135. Yamashita J. Echinostome. *Progress of Med Parasitol in Japan* 1964; Vol. 1: 287-313. Tokyo, Japan. Meguro Parasitological Museum.
 136. Kostadinova A, Gibsion DI, Biserkow V, Chipev N. Re-validation of *Echinostoma miyagawai* Ishii, 1932 (Digenea: Echinostomatidae) on the basis of the experimental completion of its life-cycle. *Syst Parasitol* 2000; 45: 81-108.
 137. Anazawa K. On a human case of *Echinostoma revolutum* and its infection route. *Taiwan Igakkai Zasshi* 1929; 288: 221-241.
 138. Ashford RW, Crewe W. The parasites of *Homo sapiens*. London, England. Liverpool School of Tropical Medicine. 2002.
 139. Radomyos P, Bunnag D, Harinasuta T. Report of *Episthmium caninum* (Verma, 1935) Yamaguti, 1958 (Digenea: Echinostomatidae) in man. *Southeast Asian J Trop Med Pub Health* 1985; 16: 508-511.
 140. Kostadinova A, Gibson DI. A redescription of *Uroproctepisthmium bursicola* (Creplin, 1837) n. comb. (Digenea: Echinostomatidae), and re-evaluations of the genera *Episthmium* Lühe, 1909 and *Uroproctepisthmium* Fischthal & Kuntz, 1976. *Syst Parasitol* 2001; 50: 63-67.
 141. Radomyos P, Charoenlarp P, Harinasuta T. Human *Episthmium caninum* (Digenea: Echinostomatidae) infection: report of two more cases. *J Trop Med Parasitol* 1991; 14: 48-50.
 142. Mas-Coma S, Bargues MD, Valero MA. Fascioliasis and other plant-borne trematode zoonoses. *Int J Parasitol* 2005; 35: 1255-1278.
 143. Rohela M, Jamaiah I, Menon J, Rachel J. Fasciolopsiasis: a first case report from Malaysia. *Southeast Asian J Trop Med Public Health* 2005; 36: 456-458.

144. Bunnag D, Radomyos P, Harinasuta C. Field trial on the treatment of fasciolopsiasis with praziquantel. *Southeast Asian J Trop Med Public Health* 1983; 14: 216-219.
145. Shyu LY, Lee HH, Chen ER. A preliminary study on epidemiology of fasciolopsiasis in Tainan Hsien, south Taiwan. *Chinese J Microbiol Immunol* 1984; 17: 118-120.
146. Lee HH. Survey of *Fasciolopsis buski* infection among children of Liu-ying primary school in Tainan county. *Chinese J Microbiol Immunol* 1972; 5: 110-114.
147. Muttalib MA, Islam N. *Fasciolopsis buski* in Bangladesh—a pilot study. *J Trop Med Hyg* 1975; 78: 135-137.
148. Lee SH, Chai JY, Hong ST. *Gymnophalloides seoi* n. sp. (Digenea: Gymnophallidae), the first report of human infection by a gymnophallid. *J Parasitol* 93; 79: 677-680.
149. Chai JY, Choi MH, Yu JR, Lee SH. *Gymnophalloides seoi*: a new human intestinal trematode. *Trends Parasitol* 2003; 19: 109-112.
150. Lee SH, Chai JY, Lee HJ, Hong ST, Yu JR, Sohn WM, Kho WG, Choi MH, Lim YJ. High prevalence of *Gymnophalloides seoi* infection in a village on a southwestern island of the Republic of Korea. *Am J Trop Med Hyg* 1994; 51: 281-285.
151. Chai JY, Kim IM, Seo M, Guk SM, Kim JL, Sohn WM, Lee SH. A new focus of *Heterophyes nocens*, *Pygidiopsis summa*, and other intestinal flukes in a coastal area of Muan-gun, Chollanam-do. *Korean J Parasitol* 1997; 35: 233-238.
152. Chai JY, Song TE, Han ET, Guk SM, Park YK, Choi MH, Lee SH. Two endemic foci of heterophyids and other intestinal fluke infections in southern and western coastal areas in Korea. *Korean J Parasitol* 1998; 36: 155-161.
153. Lee SH, Sohn WM, Hong SJ, Huh S, Seo M, Choi MH, Chai JY. A nationwide survey of naturally produced oysters for infection with *Gymnophalloides seoi* metacercariae. *Korean J Parasitol* 1996; 34: 107-112.
154. Chai JY, Park JH, Han ET, Shin EH, Kim JL, Hong KS, Rim HJ, Lee SH. A nationwide survey of the prevalence of human *Gymnophalloides seoi* infection on western and southern coastal islands in the Republic of Korea. *Korean J Parasitol* 2001; 39: 23-30.
155. Park JH, Kim JL, Shin EH, Guk SM, Park YK, Chai JY. A new endemic focus of *Heterophyes nocens* and other heterophyid infections in a coastal area of Gangjin-gun, Jeollanam-do. *Korean J Parasitol* 2007; 45: 33-38.
156. Guk SM, Park JH, Shin EH, Kim JL, Lin A, Chai JY. Prevalence of *Gymnophalloides seoi* infection in coastal villages of Haenam-gun and Yeongam-gun, Republic of Korea. *Korean J Parasitol* 2006; 44: 1-5.
157. Seo M, Shin DH, Guk SM, Oh CS, Lee EJ, Shin MH, Kim MJ, Lee SD, Kim YS, Yi YS, Spigelman M, Chai JY. *Gymnophalloides seoi* eggs from the stool of a 17th century female mummy found in hadong, Republic of Korea. *J Parasitol* 2008; 84: 467-472.
158. Lee SH, Chai JY. A review of *Gymnophalloides seoi* (Digenea: Gymnophallidae) and human infections in the Republic of Korea. *Korean J Parasitol* 2001; 39: 85-118.
159. Lee SH, Choi M H, Seo M, Chai JY. Oysters, *Crassostrea gigas*, as the second intermediate host of *Gymnophalloides seoi* (Gymnophallidae). *Korean J Parasitol* 1995; 33: 1-7.
160. Sohn WM, Ryang YS, Chai JY, Lee SH. Discovery of *Gymnophalloides seoi* metacercariae in oysters from islands of the West Sea known as the habitats of paleartic oystercatchers. *Korean J Parasitol* 1998; 36: 163-169.
161. Ryang YS, Yoo JC, Lee SH, Chai JY. The Palearctic oystercatcher *Haematopus ostralegus*, a natural definitive host for *Gymnophalloides seoi*. *J Parasitol* 2000; 86: 418-419.
162. Shin EH, Park JH, Guk SM, Kim JL, Chai JY. Intestinal helminth infections in feral cats and a raccoon dog on Aphaedo Island, Shinan-gun, with a special note on *Gymnophalloides seoi* infection in cats. *Korean J Parasitol* 2009; 47: 189-191.
163. Ryang YS, Yoo JC, Lee SH, Chai JY. Susceptibility of avian hosts to experimental *Gymnophalloides seoi* infection. *J Parasitol* 2001; 87: 454-456.
164. Lee SH, Park SK, Seo M, Guk SM, Choi MH, Chai JY. Susceptibility of various animals and strains of mice to *Gymnophalloides seoi* infection and the effects of immunosuppression in C3H/HeN mice. *J Parasitol* 1997; 83: 883-886.
165. Kook J, Lee SH, Chai JY. In vitro cultivation of *Gymnophalloides seoi* metacercariae (Digenea: Gymnophallidae). *Korean J Parasitol* 1997; 35: 25-29.
166. Lee SH, Kim HJ, Pyo KH, Shin EH, Chai JY. Effects of several culture conditions on in vitro growth and development of *Gymnophalloides seoi*. *J Nat Acad Sci, Korea (Nat Sci Ser)* 2009; 48: 199-212.
167. Radomyos P, Bunnag D, Harinasuta T. *Haplorchis pumilio* (Looss) infection in man in northeastern Thailand. *Southeast Asian J Trop Med Pub Health* 1983; 14: 223-227.
168. Chen HT. A study of the Haplorchinae (Looss, 1899) Poche 1926 (Trematoda: Heterophyidae). *Parasitology* 1936; 28: 40-55.
169. Faust EC, Nishigori M. The life cycles of two new species of Heterophyidae parasitic in mammals and birds. *J Parasitol* 1926; 13: 91-131.
170. Velasquez CC. Heterophyidiasis, In Steele JH Ed, *CRC Handbook Series in Zoonoses, Section C: Parasitic Zoonoses, Vol. III (Trematode Zoonoses)*. Boca Raton, USA. CRC Press 1982, 99-107.
171. Chai JY, Park JH, Han ET, Guk SM, Shin EH, Lin A, Kim JL, Sohn WM, Yong TS, Eom KS, Min DY, Hwang EH, Phommasack B, Insisiengmai B, Rim HJ. Mixed infections with *Opisthorchis viverrini* and intestinal flukes in residents of Vientiane Municipality and Saravane Province in Laos. *J Helminthol* 2005; 79: 283-289.
172. Dung DT, van De N, Waikagul J, Dalsgaard A, Chai JY, Sohn WM, Murrell KD. Fishborne zoonotic intestinal trematodes, Vietnam. *Emerg Infect Dis* 2007; 13: 1828-1833.
173. Rim HJ, Sohn WM, Yong TS, Eom KS, Chai JY, Min DY, Lee SH, Hoang EH, Phommasack B, Insisiengmay S. Fishborne trematode metacercariae detected in freshwater fish from Vientiane Municipality and Savannakhet Province, Lao PDR. *Korean J Parasitol* 2008; 6: 253-260.

174. Ujjiie N. Differentiation of the species of *Monorchotrema* from Formosa. Taiwan Igakkai Zasshi 1936b; 35: 938-946.
175. Africa CM, de Leon W, Garcia EY. Visceral complications in intestinal heterophyidiasis of man. Acta Medica Philippina 1940; Monographic Series No. 1: 1-132.
176. Chai JY, Han ET, Shin EH, Sohn WM, Yong TS, Eom KS, Min DY, Um JY, Park MS, Hoang EH, Phommasack B, Insiengmay B, Lee SH, Rim HJ. High prevalence of *Haplorchis taichui*, *Prosthodendrium molenkampi*, and other helminth infections among people in Khammouane province, Lao PDR. Korean J Parasitol 2009; 47: 243-247.
177. Belizario VY, Jr de Leon WU, Bersabe MJ, Purnomo BJK, Bangs MJ. A focus of human infection by *Haplorchis taichui* (Trematoda: Heterophyidae) in the southern Philippines. J Parasitol 2004; 90: 1165-1169.
178. Kumchoo K, Wongsawad C, Chai JY, Vanittanakom P, Rojanapaibul A. High prevalence of *Haplorchis taichui* metacercariae in cyprinoid fish from Chiang Mai Province, Thailand. Southeast Asian J Trop Med Public Health 2005; 36: 451-455.
179. Pearson JC, Ow-Yang CK. New species of *Haplorchis* from Southeast Asia, together with keys to the *Haplorchis*-group of heterophyid trematodes of the region. Southeast Asian J Trop Med Public Health 1982; 13: 35-60.
180. Katsuta I. Studies on the metacercariae of Formosan brackish water fishes. (4) A new trematode, *Monorchotrema yokogawai*, which takes *Mugil cephalus* as its intermediate host. Taiwan Igakkai Zasshi 1932; 31: 160-175.
181. Onji Y, Nishio T. On the trematodes whose intermediate host is brackish water fishes. Chiba Igaku Semmon Gakko Zasshi 1916; 81&82: 229-249.
182. Yokogawa M, Sano M, Itabashi T, Kachi S. Studies on the intestinal flukes II. Epidemiological studies on heterophyid trematodes of man in Chiba Prefecture. Jpn J Parasitol 1965; 14: 577-585.
183. Xu BK, Li DN. Rare human parasites in China. National Med J China 1979; 59: 286-290.
184. Seo BS, Hong ST, Chai JY. Studies on intestinal trematodes in Korea III. Natural human infection of *Pygidiopsis summa* and *Heterophyes heterophyes nocens*. Seoul J Med 1981; 22: 228-235.
185. Chai JY, Seo BS, Lee SH. Studies on intestinal trematodes in Korea XI. Two cases of human infection by *Heterophyes heterophyes nocens*. Korean J Parasitol 1984; 22: 37-42.
186. Park JH, Kim JL, Shin EH, Guk SM, Park YK, Chai JY. A new endemic focus of *Heterophyes nocens* and other heterophyid infections in a coastal area of Gangjin-gun, Jeollanam-do. Korean J Parasitol 2007; 45: 33-38.
187. Guk SM, Shin EH, Kim JL, Sohn WM, Hong KS, Yoon CH, Lee SH, Rim HJ, Chai JY. A survey of *Heterophyes nocens* and *Pygidiopsis summa* metacercariae in mullets and gobies along the coastal areas of the Republic of Korea. Korean J Parasitol 2007; 45: 205-211.
188. Eom KS, Son SY, Lee JS, Rim HJ. Heterophyid trematodes (*Heterophyopsis continua*, *Pygidiopsis summa* and *Heterophyes heterophyes nocens*) from domestic cats in Korea. Korean J Parasitol 1985; 23: 197-202.
189. Chai JY, Park JH, Han ET, Shin EH, Kim JL, Guk SM, Hong KS, Lee SH, Rim HJ. Prevalence of *Heterophyes nocens* and *Pygidiopsis summa* infections among residents of the western and southern coastal islands of the Republic of Korea. Am J Trop Med Hyg 2004; 71: 617-622.
190. Suzuki N, Kumazawa H, Araki T, Hara M, Morita H, Segawa T, Shina T, Yamada S, Tsumoto S. Occurrence of *Heterophyes heterophyes nocens* Onji et Nishio, 1916 in man in Kochi Prefecture. Jpn J Parasitol 1982; 31: 271-274.
191. Kino H, Oishi H, Ohno Y, Ishiguro M. An endemic human infection with *Heterophyes nocens* Onji et Nishio, 1916 at Mikkabicho, Shizuoka, Japan. Jpn J Trop Med Hyg 2002; 30: 301-304.
192. Africa CM, Garcia EY. Heterophyid trematodes of man and dog in the Philippines with descriptions of three new species. Phil J Sci 1935; 57: 253-267.
193. Yamaguti S. Studies on the helminth fauna of Japan. Part 27. Trematodes of mammals, II. Jpn J Med Sci (VI. Bacteriology and Parasitology) 1939; 1: 131-151.
194. Hong SJ, Chang CK, Lee DH, Woo HC. One human case of natural infection by *Heterophyopsis continua* and three other species of intestinal trematodes. Korean J Parasitol 1996; 34: 87-90.
195. Chun SK. A study on some trematodes whose intermediate hosts are brackish water fish (I) The life history of *Heterophyes continua*, the intermediate host of which is *Lateolabrax japonicus*. Bull Pusan Fish Coll 1960; 3: 40-44.
196. Sohn WM, Han GG, Kho WG, Chai JY, Lee SH. Infection status with the metacercariae of heterophyid flukes in the brackish water fish from Haenam-gun, Chollanam-do, Korea. Korean J Parasitol 1994; 32: 163-169.
197. Kim KH, Choi ES, Rim HJ. *Conger myriaster*, a new second intermediate host of *Heterophyopsis continua* (Digenea: Heterophyidae). Korean J Parasitol 1996; 34: 283-285.
198. Cho SY, Kim SI. *Plecoglossus altivelis* as a new fish intermediate host of *Heterophyopsis continua*. Korean J Parasitol 1985; 23: 173-174.
199. Hong SJ, Lee SH, Chai JY, Seo BS. Recovery rate, growth and development of *Heterophyopsis continua* in experimental chicks. Korean J Parasitol 1990; 28: 53-62.
200. Hong SJ, Chai JY, Lee SH. Surface ultrastructure of the developmental stages of *Heterophyopsis continua* (Trematoda: Heterophyidae). J Parasitol 1991; 77: 613-620.
201. Harinasuta T, Bunnag D, Radomyos P. Intestinal fluke infections. In Bailliere's Clinical Tropical Medicine and Communicable Diseases 1987; 2: 695-721.
202. Yurlova NI. Morphology and biology of trematodes *Hypoderaeum cubanicum* comb. nov. and *H. conoideum* (Echinostomatidae). Parazitologiya 1987; 21: 35-42.
203. Toledo R, Muñoz-Antolí, Pérez M, Esteban JG. Miracidial infectivity of *Hypoderaeum conoideum* (Trematoda: Echinostomatidae): differential susceptibility of two lymnaeid species. Parasitol Res 1999; 85: 212-215.
204. Yokogawa M, Harinasuta C, Charoenlarp P. *Hypoderaeum conoi-*

- deum* (Block, 1782) Diez, 1909, a common intestinal fluke of man in the north-east Thailand. *Jpn J Parasitol* 1965; 14: 148-153.
205. Shimalov VV, Shimalov VT. Human parasitizing echinostomatata (Trematoda, Fasciolidae) in mammals in Belorussia forest zone. *Meditinskaiia Parazitologija i Parazitamyje Bolezni* 2002; 1: 47-50.
206. Katsuta I. Studies on the metacercariae of Formosan brackish water fishes. (2) On a new species, *Metagonimus minutus* n. sp., parasitic in *Mugil cephalus*. *Taiwan Igakkai Zasshi* 1932; 31: 26-39.
207. Saito S, Chai JY, Kim KH, Lee SH, Rim HJ. *Metagonimus miyatai* sp. nov. (Digenea: Heterophyidae), a new intestinal trematode transmitted by freshwater fishes in Japan and Korea. *Korean J Parasitol* 1997; 3: 223-232.
208. Kim CH. Study on the *Metagonimus* sp. in Gum River Basin, Chungchung-nam Do, Korea. *Korean J Parasitol* 1980; 18: 215-228.
209. Kim CH, Kim NM, Lee CH, Park JS. Studies on the *Metagonimus* fluke in the Daechong Reservoir and the upper stream of Geum River, Korea. *Korean J Parasitol* 1987; 25: 69-82.
210. Chai JY, Huh S, Yu JR, Kook J, Jung KC, Park EC, Sohn WM, Hong ST, Lee SH. An epidemiological study of metagonimiasis along the upper reaches of the Namhan River. *Korean J Parasitol* 1993; 31: 99-108.
211. Park MS, Kim SW, Yang YS, Park CH, Lee WT, Kim CU, Lee EM, Lee SU, Huh S. Intestinal parasite infections in the inhabitants along the Hantan River, Chorwon-gun. *Korean J Parasitol* 1993; 31: 375-378.
212. Ahn YK. Intestinal fluke of genus *Metagonimus* and their second intermediate hosts in Kangwon-do. *Korean J Parasitol* 1993; 31: 331-340.
213. Kino H, Suzuki Y, Oishi H, Suzuki S, Yamagiwa S, Ishiguro M. Geographical distribution of *Metagonimus yokogawai*, and *M. miyatai* in Shizuoka Prefecture, Japan, and their site preferences in the sweetfish, *Plecoglossus altivelis*, and hamsters. *Parasitol Int* 2006; 55: 201-206.
214. Shimazu T. Life cycle and morphology of *Metagonimus miyatai* (Digenea: Heterophyidae). *Parasitol Int* 2002; 51: 271-280.
215. Guk SM, Park JY, Seo M, Han ET, Kim JL, Chai JY. Susceptibility of inbred mouse strains to infection with three species of *Metagonimus* prevalent in the Republic of Korea. *J Parasitol* 2005; 91: 12-16.
216. Suzuki S. Yokogawa's *Metagonimus*. List of publications on special animals in Okayama Prefecture. Okayama Prefecture Report 1930, p 146-148.
217. Saito S. On the validity of *Metagonimus* spp. Proceedings of Japanese Parasite Taxonomy and Morphology Conference 1984; 2: 1-6.
218. Chai JY, Sohn WM, Kim MH, Hong ST, Lee SH. Three morphological types of the genus *Metagonimus* encysted in the dace, *Tribolodon taczanowskii*, caught from the Sumjin River. *Korean J Parasitol* 1991; 29: 217-225.
219. Chun SK. A study on the metacercaria of *Metagonimus takahashii* and *Exorchis oviformis* from *Carassius carassius*. *Bull Pusan Fish Coll* 1960; 3: 31-39.
220. Cho SY, Kang SY, Lee JB. Metagonimiasis in Korea. Session V: Intestinal fluke and *Schistosoma japonicum*. *Arzneimittel-Forschung/Drug Research* 1984; 34: 1211-1213.
221. Kim DG, Kim TS, Cho SH, Song HJ, Sohn WM. Heterophyid metacercarial infections in brackish water fishes from Jinju-man (Bay), Kyongsangnam-do, Korea. *Korean J Parasitol* 2006; 44: 7-13.
222. Seo BS, Lee SH, Cho SY, Chai JY, Hong ST, Han IS, Shon JS, Cho BH, Ahn SR, Lee SK, Chung SC, Kang KS, Shim HS, Hwang IS. An epidemiologic study on clonorchiasis and metagonimiasis in riverside areas in Korea. *Korean J Parasitol* 1981; 19: 137-150.
223. Song CY, Lee SH, Jeon SR. Studies on the intestinal fluke, *Metagonimus yokogawai* Katsurada, 1912 in Korea IV. Geographical distribution of sweetfish and infection status with *Metagonimus* metacercaria in south-eastern area of Korea. *Korean J Parasitol* 1985; 23: 123-138.
224. Chai JY, Cho SY, Seo BS. Study on *Metagonimus yokogawai* (Katsurada, 1912) in Korea. IV. An epidemiological investigation along Tamjin River Basin, South Cholla Do, Korea. *Korean J Parasitol* 1977; 15: 115-120.
225. Chai JY, Han ET, Park YK, Guk SM, Kim JL, Lee SH. High endemicity of *Metagonimus yokogawai* infection among residents of Samchok-shi, Kangwon-do. *Korean J Parasitol* 2000; 38: 33-36.
226. Lee JJ, Kim HJ, Kim MJ, Yi Lee JW, Jung BK, Lee JY, Shin EH, Kim JL, Chai JY. Decrease of *Metagonimus yokogawai* endemicity along the Tamjin River basin. *Korean J Parasitol* 2008; 46: 289-291.
227. Ito J, Mochizuki H, Ohno Y, Ishiguro M. On the prevalence of *Metagonimus* sp. among the inhabitants at Hamamatsu basin in Shizuoka Prefecture, Japan. *Jpn J Parasitol* 1991; 40: 274-278.
228. Ahn YK. Lateolaborax japonicus, a role of the second intermediate host of *Metagonimus yokogawai*. *New Med J (Korea)* 1983; 26: 135-139.
229. Huh S, Sohn WM, Chai JY. Intestinal parasites of cats purchased in Seoul. *Korean J Parasitol* 1993; 31: 371-373.
230. Seo BS. *Fibricola seoulensis* Seo, Rim and Lee, 1964 (Trematoda) and fibricoliasis in man. *Seoul J Med* 1990; 31: 61-96.
231. Quan F, Jiang T, Ma H, Cui C, Sun L. The first discovery of *Fibricola seoulensis* in China. *J Yanbian Med Coll* 1995; 18: 17-20.
232. Seo BS, Lee SH, Hong ST, Hong SJ, Kim CY, Lee HY. Studies on intestinal trematodes in Korea V. A human case infected by *Fibricola seoulensis* (Trematoda: Diplostomatidae). *Korean J Parasitol* 1982; 20: 93-99.
233. Hong ST, Cho TK, Hong SJ, Chai JY, Lee SH, Seo BS. Fifteen human cases of *Fibricola seoulensis* infection in Korea. *Korean J Parasitol* 1984; 22: 61-65.
234. Hong ST, Chai JY, Lee SH. Ten human cases of *Fibricola seoulensis* infection and mixed one with *Stellantchasmus* and *Metagonimus*. *Korean J Parasitol* 1986; 24: 94-96.
235. Huh S, Lee SU, Huh SC. A follow-up examination of intestinal parasitic infections of the Army soldiers in Whachon-gun, Korea.

- Korean J Parasitol 1994; 32: 61-63.
236. Seo BS, Lee SH, Chai JY, Hong SJ, Hong ST. The life cycle and larval development of *Fibricola seoulensis* (Trematoda: Diplostomatidae). Korean J Parasitol 1988; 26: 179-188.
 237. Chung PR, Jung Y, Kim DS. *Segmentina (Polypylis) hemisphaerula* (Gastropoda: Planorbidae): a new molluscan intermediate host of a human intestinal fluke *Neodiplostomum seoulense* (Trematoda: Diplostomatidae) in Korea. J Parasitol 1996; 82: 336-338.
 238. Chai JY, Park JH, Guk SM, Kim JL, Kim HJ, Kim WH, Shin EH, Klein TA, Kim HC, Chong ST, Song JW, Baek LJ. *Apodemus agrarius* as a new definitive host for *Neodiplostomum seoulense*. Korean J Parasitol 2007; 45: 157-161.
 239. Manning GS, Diggs CL, Viyanant V, Lertprasert P, Watanasirmit K. Preliminary report on *Phaneropsolus bonnei* Lie Kian Joe, 1951. A newly discovered human intestinal fluke from northeastern Thailand. J Med Assoc Thailand 1970; 53: 173-178.
 240. Manning GS, Anluchai T, Nganpanya B, Promano R, Kanhaviang K. Redescription of the intestinal fluke *Phaneropsolus bonnei* Lie Kian Joe, 1951 (Trematoda: Lecithodendriidae). Southeast Asian J Trop Med Pub Health 1970; 1: 492-495.
 241. Manning GS, Lertprasert P, Watanasirmit K, Chetty C. A description of newly discovered intestinal parasites endemic to northeastern Thailand. J Med Assoc Thailand 1971; 54: 466-474.
 242. Kaewkes S, Elkins DB, Sithithaworn P, Haswell-Elkins MR. Comparative studies on the morphology of the eggs of *Opisthorchis viverrini* and lecithodendriid trematodes. Southeast Asian J Trop Med Public Health 1991; 22: 623-630.
 243. Tesana S, Srisawangwonk, Kaewkes S, Sithithaworn P, Kanla P, Arunyanart C. Eggshell morphology of the small eggs of human trematodes in Thailand. Southeast Asian J Trop Med Public Health 1991; 22: 631-636.
 244. Manning GS, Lertprasert P. Studies on the life cycle of *Phaneropsolus bonnei* and *Prosthodendrium molenkampii* in Thailand. Ann Trop Med Parasit 1973; 67: 361-365.
 245. Kaewkes S, Elkins DB, Haswell-Elkins MR, Sithithaworn P. *Phaneropsolus spinicirrus* n. sp. (Digenea: Lecithodendriidae), a human parasite in Thailand. J Parasitol 1991; 77: 514-516.
 246. Radomyos P, Bunnag D, Harinasuta C. A new intestinal fluke, *Plagiorchis harinasutai*, n. sp. Southeast Asian J Trop Med Public Health 1989; 20: 101-107.
 247. Sandground JH. *Plagiorchis javensis* n. sp. a new trematode parasitic in man. Rev Med Trop Parasitol (Habana) 1940; 6: 207-211.
 248. Tanabe H. A contribution to the study of the life cycle of digenetic trematodes. A study of a new species *Lepoderma muris* n. sp. Okayama Igakkai Zasshi 1922; 385: 47-58.
 249. McMullen DB. An experimental infection of *Plagiorchis muris* in man. J Parasitol 1937; 23: 113-115.
 250. Asada JI, Otagaki H, Morita M, Takeuchi T, Sakai Y, Konoshi T, Okahashi K. A case report on the human infection with *Plagiorchis muris* Tanabe, 1922 in Japan. Jpn J Parasitol 1962; 11: 512-516.
 251. Hong SJ, Woo HC, Chai JY. A human case of *Plagiorchis muris* (Tanabe, 1922: Digenea) infection in the Republic of Korea: Freshwater fish as a possible source of infection. J Parasitol 1996; 82: 647-649.
 252. Chai JY, Park JH, Guk SM, Kim JL, Kim HJ, Kim WH, Shin EH, Klein TA, Kim HC, Chong ST, Song JW, Baek LJ. *Plagiorchis muris* infection in *Apodemus agrarius* from northern Gyeonggi-do (Province) near the demilitarized zone. Korean J Parasitol 2007; 45: 153-156.
 253. Sogandares-Bernal F. Four trematodes from Korean bats with description of three new species. J Parasitol 1956; 42: 200-206.
 254. Guk SM, Kim JL, Park JH, Chai JY. A human case of *Plagiorchis vespertilionis* (Digenea: Plagiorchiidae) infection in the Republic of Korea. J Parasitol 2007; 93: 1225-1227.
 255. Macy RW. The life cycle of *Plagiorchis vespertilionis* Parorchis. N. SSP. (Trematoda: Plagiorchiidae), and observations on the effects of light on the emergence of the cercaria. J Parasitol 1960; 46: 337-345.
 256. Africa CM, Garcia EY. Heterophyid trematodes of man and dog in the Philippines with descriptions of three new species. Philippine J Sci 1935; 57: 253-267.
 257. Velasquez CC. Life cycle of *Procerovum calderoni* (Africa and Garcia, 1935), Price, 1940 (Trematoda: Digenea: Heterophyidae). J Parasitol 1973; 59: 813-816.
 258. Velasquez CC. Observations on some Heterophyidae (Trematoda: Digenea) encysted in Philippine fishes. J Parasitol 1973; 59: 77-84.
 259. Aokage K. Studies on the trematode parasites of brackish water fishes in Chugoku coast of Setonaikai. Tokyo Iji Shinshi 1956; 73: 217-224.
 260. Umadevi K, Madhavi R. Observation on the morphology and life-cycle of *Procerovum varium* (Onji & Nishio, 1916) (Trematoda: Heterophyidae). Syst Parasitol 2000; 46: 215-225.
 261. Kifune T, Takao Y. Description of *Psilorchis hominis* sp. nov. from man (Trematoda: Echinostomatoidea: Psilostomidae). Jpn J Parasitol 1973; 22: 111-115.
 262. Chai JY, Seo BS, Lee SH, Hong ST. Growth and development of *Pygidiopsis summa* in rats and mice with a supplementary note on its morphological characters. Korean J Parasitol 1986; 24: 55-62.
 263. Takahashi S. On the eggs of *Stellantchasmus falcatus* and *Pygidiopsis summa* found in human stools. Okayama Igakkai Zasshi 1929; 41: 1502-1513.
 264. Chun SK. A study on some trematodes whose intermediate hosts are brackish water fish (II) The life history of *Pygidiopsis summa*, the intermediate host of which is *Mugil cephalus*. Bull Pusan Fish Coll 1963; 5: 1-5.
 265. Seo BS, Hong ST, Chai JY, Cho SY. Studies on intestinal trematodes in Korea IV. Geographical distribution of *Pygidiopsis* and *Heterophyes* metacercariae. Seoul J Med 1981; 22: 236-242.
 266. Kagei N, Oshima T, Ishikawa K, Kihata M. Two cases of human infection with *Stellantchasmus falcatus* Onji et Nishio, 1915 (Heterophyidae) in Kochi Prefecture. Jpn J Parasitol 1964; 13: 472-

- 478.
267. Radomyos P, Charoenlarp P, Radomyos B, Tungtrongchitr A. Two human cases of *Stellantchasmus falcatus* (Trematoda, Heterophyidae) infection in Northeastern Thailand. *Jpn J Parasitol* 1990; 39: 7-11.
268. Seo BS, Lee SH, Chai JY, Hong SJ. Studies on intestinal trematodes in Korea XII. Two cases of human infection by *Stellantchasmus falcatus*. *Korean J Parasitol* 1984; 22: 43-50.
269. Noda K. The larval development of *Stellantchasmus falcatus* (Trematoda; Heterophyidae) in the first intermediate host. *J Parasitol* 1959; 45: 635-642.
270. Chai JY, Sohn WM. Identification of *Stellantchasmus falcatus* metacercariae encysted in mullets in Korea. *Korean J Parasitol* 1988; 26: 65-68.
271. Wongsawad C, Chariyahpongpun P, Namue C. Experimental host of *Stellantchasmus falcatus*. *Southeast Asian J Trop Med Public Health* 1998; 29: 406-409.
272. Sripalwit P, Wongsawad C, Chai JY, Anuntalabhochai S, Rojana-paibul A. Investigation of *Stellantchasmus falcatus* metacercariae in half-beaked fish, *Dermogenus pusillus* from four districts of Chiang Mai Province, Thailand. *Southeast Asian J Trop Med Public Health* 2003; 34: 281-285.
273. Saenphet S, Wongsawad C, Saenphet K, Chai JY. Susceptibility of rodents to *Stellantchasmus falcatus* infection. *Southeast Asian J Trop Med Public Health* 2003; 34: 123-127.
274. Katsuta I. Studies on the metacercariae of Formosan brackish water fishes. (1) On a new species, *Stellantchamus formosanus* n. sp., parasitic in *Mugil cephalus*. *Taiwan Igakkai Zasshi* 1931; 30: 1404-1417.
275. Witenberg G. Studies on the trematode-family Heterophyidae. *Ann Trop Med Parasitol* 1929; 23: 131-268.
276. Katsuta I. Studies on the metacercariae of Formosan brackish water fishes. (5) On a new species, *Stellantchasmus amplicaealis* n. sp., parasitic in *Mugil cephalus*. *Taiwan Igakkai Zasshi* 1932; 31: 457-471.
277. Chai JY, Hong SJ, Lee SH, Seo BS. *Stictodora* sp. (Trematoda: Heterophyidae) recovered from a man in Korea. *Korean J Parasitol* 1988; 26: 127-132.
278. Cohn WM, Chai JY, Lee SH. *Stictodora fuscatum* (Heterophyidae) metacercariae encysted in gobies, *Acanthogobius flavimanus*. *Korean J Parasitol* 1994; 32: 143-148.
279. Chai JY, Park SK, Hong SJ, Choi MH, Lee SH. Identification of *Stictodora lari* (Heterophyidae) metacercariae encysted in the brackish water fish, *Acanthogobius flavimanus*. *Korean J Parasitol* 1989; 27: 253-259.
280. Chai JY, Han ET, Park YK, Guk SM, Park JH, Lee SH. *Stictodora lari* (Digenea: Heterophyidae): the discovery of the first human infections. *J Parasitol* 2002; 88: 627-629.
281. Howell MJ. The resistance of cysts of *Stictodora lari* (Trematoda: Heterophyidae) to encapsulation by cells of the fish host. *Int J Parasitol* 1973; 3: 653-659.
282. Marcilla A. Echinotomes: genomes and proteomics. In Fried B, Toledo R eds, *The Biology of Echinostomes*. New York, USA: Springer. 2009, p 207-228.
283. Nowak TS, Loker ES. *Echinostoma paraensei*: Differential gene transcription in the sporocyst stage. *Exp Parasitol* 2005; 109: 94-105.
284. Morgan JA, Blair D. Nuclear rDNA ITS sequence variation in the trematode genus *Echinostoma*: An aid to establishing relationships within the 37-collar-spine group. *Parasitology* 1995; 111: 609-615.
285. Morgan JA, Blair D. Mitochondrial ND1 gene sequences used to identify echinostome isolates from Australia and New Zealand. *Int J Parasitol* 1998; 28: 493-502.
286. Taft JL, Fried B. Oxygen consumption in adult *Echinostoma revolutum*. *Exp Parasitol* 1968; 23: 183-186.
287. Heyneman D, Faulk WP, Fudenberg HH. *Echinostoma lindoense*: larval antigens from the snail intermediate host, *Biomphalaria glabrata*. *Exp Parasitol* 1971; 29: 480-492.
288. Fujino T, Takahashi Y, Fried B. A comparison of *Echinostoma trivolvis* and *E. caproni* using random amplified polymorphic DNA analysis. *J Helminthol* 1995; 69: 263-264.
289. Bernal D, Carpena I, Espert AM, De la Rubia JE, Esteban JC, Toledo R, Marcilla A. Identification of proteins in excretory/secretory extracts of *Echinostoma friedi* (Trematoda) from chronic and acute infections. *Proteomics* 2006; 6: 2835-2843.
290. Le TH, Nguyen VD, Phan BU, Blair D, McManus DP. Case report: unusual presentation of *Fasciolopsis buski* in a Vietnamese child. *Trans R Soc Trop Med Hyg* 2004; 98: 193-194.
291. Prasad PK, Tandon V, Chatterjee A, Bandyopadhyay S. PCR-based determination of internal transcribed spacer (ITS) regions of ribosomal DNA of giant intestinal fluke, *Fasciolopsis buski* (Lankester, 1857) Looss, 1899. *Parasitol Res* 2007; 101: 1581-1587.
292. Guk SM, Lee KD, Chai JY. Toll-like receptor 2 and muc2 expression on human intestinal epithelial cells by *Gymnophalloides seoi* adult antigen. *J Parasitol* 2009; 8: 1.
293. Leung TL, Poulin R, Keeney DB. Ten polymorphic microsatellite loci for the trematode *Gymnophallus* sp. *Mol Ecol Notes* 2007; 7: 1039-1041.
294. Leung TL, Poulin R, Keeney DB. Accumulation of diverse parasite genotypes within the bivalve second intermediate host of the digenean *Gymnophallus* sp. *Int J Parasitol* 2009; 39: 327-331.
295. Choi MH, Chai JY, Lee SH. Purification and characterization of a 16-kDa cysteine proteinase of *Gymnophalloides seoi* (Gymnophalloidae) metacercariae. *J Parasitol* 1998; 84: 350-355.
296. Choi MH, Park WJ, Park YK, Chai JY, Lee SH. Isolation and characterization of a 40 kDa cysteine protease from *Gymnophalloides seoi* adult worms. *Korean J Parasitol* 1998; 36: 133-141.
297. Lee SC, Chung YB, Kong Y, Kang SY, Cho SY. Antigenic protein fractions of *Metagonimus yokogawai* reacting with patient sera. *Korean J Parasitol* 1993; 31: 43-48.
298. Yu JR, Chung JS, Huh S, Lee SH, Chai JY. PCR-RFLP patterns of three kinds of *Metagonimus* in Korea. *Korean J Parasitol* 1997; 35: 271-276.
299. Yu JR, Chung JS, Chai JY. Different RAPD pattern between *Meta-*

- gonimus yokogawai* and *Metagonimus* Miyata type. Korean J Parasitol 1997; 35: 295-298.
300. Yu JR, Chai JY. Chapter 59. *Metagonimus*. In Liu D ed, Molecular Detection of Foodborne Pathogens. Boca Raton, Florida, USA. CRC Press (Taylor & Francis Group). 2010, p 805-811.
 301. Lee SU, Huh S, Park GM, Chai JY. A cytogenetic study on human intestinal trematodes of the genus *Metagonimus* (Digenea: Heterophyidae) in Korea. Korean J Parasitol 1999; 37: 237-241.
 302. Yang HJ, Guk SM, Han ET, Chai JY. Molecular differentiation of three species of *Metagonimus* by simple sequence repeat anchored polymerase chain reaction (SSR-PCR) amplification. J Parasitol 2000; 86: 1170-1172.
 303. Lee SU, Huh S, Sohn WM, Chai JY. Sequence comparisons of 28S ribosomal DNA and mitochondrial cytochrome C oxidase subunit I of *Metagonimus yokogawai*, *M. takahashii* and *M. miyatai*. Korean J Parasitol 2004; 42: 129-135.
 304. Dzikowski R, Levy MG, Poore MF, Flowers JR, Paperna I. Use of rDNA polymorphism for identification of Heterophyidae infecting freshwater fishes. Dis Aquat Organ 2004; 59: 35-41.
 305. Wongsawad C, Wongsawad P, Chai JY, Paratasilpin T, Anuntalabhochai S. Dna quantities and qualities from various stages of some trematodes using optical and HAT-RAPD methods. South-east Asian J Trop Med Public Health 2006; 3: 62-68.
 306. Wongsawad C, Wongsawad P, Chai JY, Anuntalabhochai S. *Haplorchis taichui*, Witenberg, 1930: development of a HAT-RAPD marker for the detection of minute intestinal fluke infection. Exp Parasitol 2009; 123: 158-161.
 307. Chai JY, Shin EH. *Neodiplostomum leei* n. sp. (Digenea: Neodiplostomidae) from chicks infected with metacercariae from the grass snake, *Rhabdophis tigrina*. J Parasitol 2002; 88: 1181-1186.
 308. Shin EH, Kim JL, Chai JY. A new neodiplostomid (Digenea) from the intestine of chicks infected with metacercariae from the grass snake *Rhabdophis tigrina*. J Parasitol 2008; 94: 1379-1384.
 309. Choi MH, Choe SC, Lee SH. A 54 kDa cysteine protease purified from the crude extract of *Neodiplostomum seoulense* adult worms. Korean J Parasitol 1999; 37: 39-46.
 310. Kim TY, Han KY, Shin EH, Chai JY. Antigenic properties of cystatin-binding cysteine proteinases from *Neodiplostomum seoulense*. J Parasitol 2008; 94: 654-658.
 311. Toledo R, Esteban JG, Fried B. Immunology and pathology of intestinal trematodes in their definitive hosts. Adv Parasitol 2006; 63: 285-365.
 312. Fried B, Huffman JE, Weiss PM. Single and multiple worm infections of *Echinostoma caproni* (Trematoda) in the golden hamster. J Helminthol 1990; 64: 75-78.
 313. Reddy DB, Ranganaykamma I, Venkataratnam D. *Artyfechinostomum mehrai* infestation in man. J Trop Med Hyg 1964; 67: 58-59.
 314. Kaul BK, Singhal GD, Pillai PN. *Artyfechinostomum mehrai* infestation with bowel perforation. J Indian Med Assoc 1974; 16: 263-265.
 315. Bindseil E, Christensen NO. Thymus-independent crypt hyperplasia and villous atrophy in the small intestine of mice infected with the trematode *Echinostoma revolutum*. Parasitology 1984; 88: 431-438.
 316. Huffman JE, Michos C, Fried B. Clinical and pathological effects of *Echinostoma revolutum* (Digenea: Echinostomatidae) in the golden hamster, *Mesocricetus auratus*. Parasitology 1986; 93: 505-515.
 317. Lee SH, Noh TY, Sohn WM, Kho WG, Hong ST, Chai JY. Chronological observation of intestinal lesions of rats experimentally infected with *Echinostoma hortense*. Korean J Parasitol 1990; 28: 45-52.
 318. Kim S, Fried B. Pathological effects of *Echinostoma caproni* (Trematoda) in the domestic chick. J Helminthol 1989; 63: 227-230.
 319. Fujino T, Fried B, Tada I. The expulsion of *Echinostoma trivolvis*: worm kinetics and intestinal cytopathology in conventional and congenitally athymic BALB/c mice. Parasitology 1993; 106: 297-304.
 320. Marty AM, Andersen EM. Fasciolopsiasis and other intestinal trematodiases. Pathol Infect Dis 2000; 1: 93-105.
 321. Bhattacharjee HK, Yadav D, Bagga D. Fasciolopsiasis presenting as intestinal perforation: A case report. Trop Gastroenterol 2009; 30: 40-41.
 322. Chai JY, Lee HS, Hong SJ, Yoo JH, Guk SM, Seo M, Choi MH, Lee SH. Intestinal histopathology and in situ postures of *Gymnophalloides seoi* in experimentally infected mice. Korean J Parasitol 2001; 39: 31-41.
 323. Seo M, Chun H, Ahn G, Jang KT, Guk SM, Chai JY. A case of colonic lymphoid tissue invasion by *Gymnophalloides seoi* in a Korean man. Korean J Parasitol 2006; 44: 87-89.
 324. Lee SH, Chai JY, Seo M, Choi MH, Kim DC, Lee SK. Two cases of *Gymnophalloides seoi* infection accompanied by diabetes mellitus. Korean J Parasitol 1995; 33: 61-64.
 325. Chai JY. Study on *Metagonimus yokogawai* (Katsurada, 1912) in Korea V. Intestinal pathology in experimentally infected albino rats. Seoul J Med 1979; 20: 104-117.
 326. Lee JB, Chi JG, Lee SK, Cho SY. Study on the pathology of metagonimiasis in experimentally infected cat intestine. Korean J Parasitol 1981; 19: 109-129.
 327. Kang SY, Cho SY, Chai JY, Lee JB, Jang DH. A study on intestinal lesions of experimentally reinfected dogs with *Metagonimus yokogawai*. Korean J Parasitol 1983; 21: 53-73.
 328. Seo BS, Cheong SK, Chai JY, Lee SH, Lee JB. Histopathology of small intestine of rats and mice experimentally infected with *Pygidioopsis summa*. Seoul J Med 1986; 27: 125-134.
 329. Hong SJ, Han JH, Park CK, Kang SY. Intestinal pathologic findings at early stage infection by *Centrocestus armatus* in albino rats. Korean J Parasitol 1997; 35: 135-138.
 330. Chi JG, Kim CW, Kim JR, Hong ST, Lee SH. Intestinal pathology in human metagonimiasis with ultrastructural observations of parasites. J Korean Med Sci 1988; 3: 171-177.
 331. Rho IH, Kim SI, Kang SY, Cho SY. Observation on the pathogenesis of villous changes in early phase of experimental metagonimiasis. Chung-Ang J Med 1984; 9: 67-77.

332. Jang YK, Kang SY, Kim SI, Cho SY. In situ posture of anterior body of *Metagonimus yokogawai* in experimentally infected dog. *Korean J Parasitol* 1985; 23: 203-213.
333. Chai JY, Kim J, Lee SH. Invasion of *Metagonimus yokogawai* into the submucosal layer of the small intestine of immunosuppressed mice. *Korean J Parasitol* 1995; 33: 313-321.
334. Chai JY, Seo BS, Lee SH. Study on *Metagonimus yokogawai* (Katsurada, 1912) in Korea VII. Susceptibility of various strains of mice to *Metagonimus* infection and effect of prednisolone. *Korean J Parasitol* 1984; 22: 153-160.
335. Hong ST, Yu JR, Myong NH, Chai JY, Lee SH. Activities of brush border membrane bound enzymes of the small intestine in *Metagonimus yokogawai* infection in mice. *Korean J Parasitol* 1990; 29: 9-20.
336. Cho SY, Kim SI, Earm YE, Ho WK. A preliminary observation on watery content of small intestine in *Metagonimus yokogawai* infected dog. *Korean J Parasitol* 1985; 23: 175-177.
337. Yu JR, Myong N, Chai JY. Expression patterns of proliferating cell nuclear antigen in the small intestine of mice infected with *Metagonimus yokogawai* and *Metagonimus Miyata* type. *Korean J Parasitol* 1997; 35: 239-244.
338. Zhang SL, Fan GH. Case report: brain abscess caused by heterophyid eggs. *Chinese J Parasitol Parasit Dis* 1990; 8: 178.
339. Lee SH, Yoo BH, Hong ST, Chai JY, Seo BS, Chi JG. A histopathological study on the intestine of mice and rats experimentally infected by *Fibricola seoulensis*. *Korean J Parasitol* 1985; 23: 58-72.
340. Huh S, Chai JY, Hong ST, Lee SH. Clinical and histopathological findings in mice heavily infected with *Fibricola seoulensis*. *Korean J Parasitol* 1988; 26: 45-53.
341. Kook J, Nawa Y, Lee SH, Chai JY. Pathogenicity and lethality of a minute intestinal fluke, *Neodiplostomum seoulense*, to various strains of mice. *J Parasitol* 1998; 84: 1178-1183.
342. Chai JY, Shin EH, Han ET, Guk SM, Choi MH, Lee SH. Genetic difference in susceptibility and fatality of three strains of mice experimentally infected with *Neodiplostomum seoulense*. *J Parasitol* 2000b; 86: 1140-1144.
343. Huh S, Lee SH, Seo BS. Histochemical findings of the tribocytic organ and tegument of *Fibricola seoulensis*. *Korean J Parasitol* 1990; 28: 155-160.
344. Cho YK, Ryang YS, Kim IS, Park SK, Im JA, Lee KJ. 2007. Differential immune profiles following experimental *Echinostoma hortense* infection in BALB/c and C3H/HeN mice. *Parasitol Res* 100: 1053-1061.
345. Ryang YS, Yang EJ, Kim JL, Lee KJ, Sung HJ, Kim JB, Kim IS. Immune response and inhibitory effect of ketotifen on the BALB/c and C3H/HeN mice infected with *Echinostoma hortense*. *Parasitol Res* 2007; 101: 1103-1110.
346. Fujino T, Fried B. *Echinostoma caproni* and *E. trivolvis* alter the binding of glycoconjugates in the intestinal mucosa of C3H mice as determined by lectin histochemistry. *J Helminthol* 1993; 67: 179-188.
347. Weinstein MS, Fried B. The expulsion of *Echinostoma trivolvis* and retention of *Echinostoma caproni* in the ICR mouse: pathological effects. *Int J Parasitol* 1991; 21: 255-257.
348. Mabus J, Huffman JE, Fried B. Humoral and cellular response to infection with *Echinostoma revolutum* in the golden hamster, *Mesocricetus auratus*. *J Helminthol* 1988; 62: 127-132.
349. Toledo R, Espert A, Carpena I, Muñoz-Antoli C, Esteban JG. An experimental study of the reproductive success of *Echinostoma friedi* (Trematoda : Echinostomatidae) in the golden hamster. *Parasitology* 2003; 126: 433-441.
350. Toledo R, Espert A, Carpena I, Muñoz-Antoli C, Fried B, Esteban JG. The comparative development of *Echinostoma caproni* (Trematoda: Echinostomatidae) adults in experimentally infected hamsters and rats. *Parasitol Res* 2004; 93: 439-444.
351. Fujino T, Fried B, Ichikawa H, Tada I. Rapid expulsion of the intestinal trematodes *Echinostoma trivolvis* and *E. caproni* from C3H mice by trapping with increased goblet cell mucins. *Int J Parasitol* 1996; 26: 319-324.
352. Kim I, Im JA, Lee KJ, Ryang YS. Mucosal mast cell responses in the small intestine of rats infected with *Echinostoma hortense*. *Korean J Parasitol* 2000; 38: 139-143.
353. Fujino T, Fried B, Tada I. The expulsion of *Echinostoma trivolvis*: Worm kinetics and intestinal cytopathology in conventional and congenitally athymic BALB/c mice. *Parasitology* 1993; 106: 297-304.
354. Fujino T, Ichikawa H, Fried B, Fukuda K. The expulsion of *Echinostoma trivolvis*: Suppressive effects of dexamethasone on goblet cell hyperplasia and worm rejection in C3H/HeN mice. *Parasite* 1996; 3: 283-289.
355. Fujino T, Ichikawa H, Fried B, Fukuda K. The expulsion of *Echinostoma trivolvis*: Worm kinetics and intestinal reactions in C3H/HeN mice treated with dexamethasone. *J Helminthol* 1997; 71: 257-259.
356. McMaster RP, Huffman JE, Fried B. The effect of dexamethasone on the course of *Echinostoma caproni* and *E. trivolvis* infections in the golden hamster (*Mesocricetus auratus*). *Parasitol Res* 1995; 81: 518-521.
357. Fujino T, Yamada M, Ichikawa H, Fried B, Arizono N, Tada I. Rapid expulsion of the intestinal trematodes *Echinostoma trivolvis* and *E. caproni* from C3H/HeN mice after infection with *Nippostrongylus brasiliensis*. *Parasitol Res* 1996; 82: 577-579.
358. Frazer BA, Fried B, Fujino T, Sleckman BP. Host-parasite relationships between *Echinostoma caproni* and RAG-2-deficient mice. *Parasitol Res.* 1999; 85: 337-342.
359. Muñoz-Antoli C, Sotillo J, Monteagudo C, Fried B, Marcilla A, Toledo R. Development and pathology of *Echinostoma caproni* in experimentally infected mice. *J Parasitol* 2007; 93: 854-859.
360. Fujino T, Fried B, Hosier DW. The expulsion of *Echinostoma trivolvis* (Trematoda) from ICR mice: Extension/retraction mechanisms and ultrastructure of the collar spines. *Parasitol Res* 1994; 80: 581-587.
361. Agger MK, Simonsen PE, Vennervald BJ. The antibody response in serum, intestinal wall and intestinal lumen of NMRI mice injected with *Echinostoma caproni*. *J Helminthol* 1993; 67: 169-

- 178.
362. Sotillo J, Muñoz-Antoli C, Macilla A, Fried B, Esteban JG, Toledo R. *Echinostoma caproni*: Kinetics of IgM, IgA and IgG subclasses in the serum and intestine of experimentally infected rats and mice. *Exp Parasitol* 2007; 116: 390-398.
 363. Simonsen PE, Andersen BJ. *Echinostoma revolutum* in mice: Dynamics of the antibody attack to the surface of an intestinal trematode. *Int J Parasitol* 1986; 16: 475-482.
 364. Shin EH, Lee SH, Kim JL, Chai JY. T-helper-1 and T-helper-2 immune responses in mice infected with the intestinal fluke *Neodiplostomum seoulense*: their possible roles in worm expulsion and host fatality. *J Parasitol* 2007; 93: 1036-1045.
 365. Brunet IR, Joseph S, Dunne DW, Fried B. Immune responses during the acute stage of infection with the intestinal trematode *Echinostoma caproni*. *Parasitology* 2000; 120: 565-571.
 366. Toledo R, Fried B. Echinostomes as experimental models for interactions between adult parasites and vertebrate hosts. *Trends Parasitol* 2005; 21: 251-254.
 367. Chai JY, Chung WJ, Kook J, Seo M, Park YK, Guk SM, Choi MH, Lee SH. Growth and development of *Gymnophalloides seoi* in immunocompetent and immunosuppressed C3H/HeN mice. *Korean J Parasitol* 1999; 37: 21-26.
 368. Seo M, Guk SM, Han ET, Chai JY. Role of intestinal goblet cells in the expulsion of *Gymnophalloides seoi* from mice. *J Parasitol* 2003; 89: 1080-1082.
 369. Guk SM, Lee JH, Kim HJ, Kim WH, Shin EH, Chai JY. CD4+ T-cell-dependent goblet cell proliferation and expulsion of *Gymnophalloides seoi* from the intestine of C57BL/6 mice. *J Parasitol* 2009; 95: 581-590.
 370. Chai JY, Yun TY, Kim J, Huh S, Choi MH, Lee SH. Chronological observation on intestinal histopathology and intraepithelial lymphocytes in the intestine of rats infected with *Metagonimus yokogawai*. *Korean J Parasitol* 1994; 32: 215-221.
 371. Chai JY, Kim TH, Kho WC, Chung SW, Hong ST, Lee SH. Mucosal mast cell responses to experimental *Metagonimus yokogawai* infection in rats. *Korean J Parasitol* 1993; 31: 129-134.
 372. Ahn H, Rim HJ, Kim SJ. Antigenic localities in the tissues of *Metagonimus yokogawai* observed by immunogold labeling method. *Korean J Parasitol* 1991; 29: 245-257.
 373. Rim HJ, Kim SJ, Yang MG. Antigenic localities in the tissues of *Metagonimus yokogawai* in the period of growth. *Korean J Parasitol* 1992; 30: 309-321.
 374. Chai JY, Kim TK, Cho WH, Seo M, Kook J, Guk SM, Lee SH. Intestinal mastocytosis and goblet cell hyperplasia in BALB/c and C3H mice infected with *Neodiplostomum seoulense*. *Korean J Parasitol* 1998; 36: 109-119.
 375. Kho WC, Chai JY, Chun CH, Lee SH. Mucosal mast cell responses to experimental *Fibricola seoulensis* infection. *Seoul J Med* 1990; 31: 191-199.
 376. Shin EH, Kim TH, Hong SJ, Park JH, Guk SM, Chai JY. Effects of anti-allergic drugs on intestinal mastocytosis and worm expulsion of rats infected with *Neodiplostomum seoulense*. *Korean J Parasitol* 2003; 41: 81-87.
 377. Han ET, Chen JH, Chai JY. Antibody responses in sera of different mouse strains experimentally infected with *Neodiplostomum seoulense*. *Korean J Parasitol* 2008; 46: 279-283.
 378. Huh S, Lee SU, Won MH, Jeong YG, Kwon YH, Choi CS. IgA response in mice infected with *Neodiplostomum seoulensis*. *Korean J Parasitol* 1995; 33: 55-60.
 379. Lee JC, Kong Y, Lee SU, Huh S. Localization of worm antigen in *Neodiplostomum seoulense* by immuno-electronmicroscopy. *Korean J Parasitol* 1997; 35: 95-104.
 380. Lee SK, Chung NS, Ko IH, Ko HI, Chai JY. Two cases of natural human infection by *Echinostoma hortense*. *Korean J Parasitol* 1986; 24: 77-81.
 381. Poland GA, Navin TR, Sarosi GA. Outbreak of parasitic gastroenteritis among travelers returning from Africa. *Arch Intern Med* 1985; 145: 2220-2221.
 382. Hamamoto T, Kawasaki H, Maejima J, Hirai K. A case of *Echinostoma hortense* infection diagnosed by the upper gastrointestinal endoscopy. *Jpn J Gastroenterol Soc* 1997; 94: 487-491.
 383. Gilman RH, Mondal G, Maksud M, Alam K, Rutherford E, Gilman JB, Khan MU. Endemic focus of *Fasciolopsis buski* infection in Bangladesh. *Am J Trop Med Hyg* 1982; 31: 796-802.
 384. Chai JY, Lee GC, Park YK, Han ET, Seo M, Kim J, Guk SM, Shin EH, Choi MH, Lee SH. Persistent endemicity of *Gymnophalloides seoi* infection in a southwestern coastal village with special reference to its egg laying capacity in the human host. *Korean J Parasitol* 2000; 38: 51-57.
 385. Seo BS, Lee HS, Chai JY, Lee SH. Intensity of *Metagonimus yokogawai* infection among inhabitants in Tamjin River basin with reference to its egg laying capacity in human host. *Seoul J Med* 1985; 26: 207-212.
 386. Chai JY, Yu JR, Lee SH, Jung HC, Song IS, Cho SY. An egg-negative patient of acute metagonimiasis diagnosed serologically by ELISA. *Seoul J Med* 1989; 30: 139-142.
 387. Chai JY, Nam HK, Kook J, Lee SH. The first discovery of an endemic focus of *Heterophyes nocens* (Heterophyidae) infection in Korea. *Korean J Parasitol* 1994; 32: 157-161.
 388. Seo BS, Lee SH, Hong ST, Hong SJ, Kim CY, Lee HY. Studies on intestinal trematodes in Korea V. A human case infected by *Fibricola seoulensis* (Trematoda: Diplostomatidae). *Korean J Parasitol* 1982; 20: 93-99.
 389. Quang TD, Duong TH, Richard-Lenoble D, Odermatt P, Khammanivong K. Emergence in humans of fascioliasis (from *Fasciola gigantica*) and intestinal distomatosis (from *Fasciolopsis buski*) in Laos. *Sante* 2008; 18: 119-124.
 390. Lee SH, Hwang SW, Chai JY, Seo BS. Comparative morphology of eggs of heterophyids and *Clonorchis sinensis* causing human infections in Korea. *Korean J Parasitol* 1984; 22: 171-180.
 391. Cho SY, Kim SI, Kang SY. Specific IgG antibody response in experimental cat metagonimiasis. *Korean J Parasitol* 1987; 25: 149-153.
 392. Datta S, Mukerjee GS, Ghosh JD. Comparative efficacy of triclabendazole, oxiclozanide and rafoxanide against *Fasciolopsis buski* in naturally infected pigs. *J Anim Health* 2004; 43: 53-56.

393. Chai JY, Han MS, Seo M, Lee SH. Effects of gamma-irradiation on the survival and development of *Gymnophalloides seoi* in C3H mice. *Korean J Parasitol* 1996; 34: 21-25.
394. Rim HJ, Chu DS, Lee JS, Joo KH, Won CY. Anthelmintic effects of various drugs against metagonimiasis. *Korean J Parasitol* 1978; 16: 117-122.
395. Lee SH, Chai JY, Hong SJ, Jun YS, Seo BS. Anthelmintic effect of praziquantel (Distocide®) against *Metagonimus yokogawai* infection. *J Korean Soc Chemother* 1984; 2: 167-171.
396. Kumchoo K, Wongsawad C, Vanittanakom P, Chai JY, Rojana-paibul A. Effect of niclosamide on the tegumental surface of *Haplorchis taichui* using scanning electron microscopy. *J Helminthol* 2007; 81: 329-337.
397. Chai JY, Kim SJ, Kook J, Lee SH. Effects of gamma-irradiation on the survival and development of *Metagonimus yokogawai* metacercariae in rats. *Korean J Parasitol* 1995; 33: 297-303.