



# Genotype and Phenotype of *Echinococcus granulosus* Derived from Wild Sheep (*Ovis orientalis*) in Iran

Ali Eslami<sup>1</sup>, Behnam Meshgi<sup>1,\*</sup>, Fatemeh Jalousian<sup>1</sup>, Shima Rahmani<sup>2</sup>, Mohammad Ali Salari<sup>2</sup>

<sup>1</sup>Department of Parasitology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran (Center of Excellence of Ecosystem and Ultrastructural Changes of Helminthes); <sup>2</sup>Under Graduate Student, School of Specialized Sciences of Veterinary Medicine, Unit of Sciences and Research, Islamic Azad University, Tehran, Iran

**Abstract:** The aim of the present study is to determine the characteristics of genotype and phenotype of *Echinococcus granulosus* derived from wild sheep and to compare them with the strains of *E. granulosus sensu stricto* (sheep-dog) and *E. granulosus* camel strain (camel-dog) in Iran. In Khojir National Park, near Tehran, Iran, a fertile hydatid cyst was recently found in the liver of a dead wild sheep (*Ovis orientalis*). The number of protoscolices (n=6,000) proved enough for an experimental infection in a dog. The characteristics of large and small hooks of metacestode were statistically determined as the *sensu stricto* strain but not the camel strain ( $P=0.5$ ). To determine *E. granulosus* genotype, 20 adult worms of this type were collected from the infected dog. The second internal transcribed spacer (ITS2) of the nuclear ribosomal DNA (rDNA) and cytochrome c oxidase 1 subunit (COX1) of the mitochondrial DNA were amplified from individual adult worm by PCR. Subsequently, the PCR product was sequenced by Sanger method. The lengths of ITS2 and COX1 sequences were 378 and 857 bp, respectively, for all the sequenced samples. The amplified DNA sequences from both ribosomal and mitochondrial genes were highly similar (99% and 98%, respectively) to that of the ovine strain in the GenBank database. The results of the present study indicate that the morpho-molecular features and characteristics of *E. granulosus* in the Iranian wild sheep are the same as those of the sheep-dog *E. granulosus sensu stricto* strain.

**Key words:** *Echinococcus granulosus*, wild sheep, genotype, phenotype, Iran

## INTRODUCTION

The pastoral cycle of echinococcosis/hydatidosis is endemic in most parts of the world, including Iran. Domestic dogs are known as the final host, while several hosts such as domestic ruminants, humans [1], and donkey in Iran [2] are identified as intermediate hosts. In the sylvatic cycle of *Echinococcus granulosus*, the wild sheep and wild boar act as the intermediate host, whereas the golden jackal [3,4] acts as the final hosts. In Iran, based on morphological characteristics of *E. granulosus*, there are 2 distinct strains, including the sheep-dog and the camel-dog strains [5]. Ungulates like wild sheep, wild goats, and gazelle ingest the egg which hatches in the intestine. Consequently, oncospheres penetrate the wall of the intestine and travel through the blood stream to become lodged in the cap-

illary beds, finally encapsulated in organs as hydatid cyst. Canids like wolves, jackals, and dogs eat hydatid cysts, and, therefore, scolices attach to their intestinal walls, grow into adult tapeworms, and produce eggs.

Based on intermediate and final hosts of *E. granulosus*, 10 distinct genotypes of G1-G10 exist in the world [6-8]. For the first time in Iran, the genotype study of 16 isolates of *E. granulosus* using mitochondrial DNA (mtDNA) markers from humans, sheep, goats, cattle, and camels revealed the existence of 2 distinct strains; the *sensu stricto* strain which infects sheep, goats, cattle, and humans, and the camel strain which infects camels [9]. The *sensu stricto* strain (sheep-dog) was recently determined in buffalos throughout Iran, except in a very limited area in its west as the buffalo-dog or G3 strain [10]. Further studies showed the possibility of interaction among the cycles of transmission.

Sheep is the most important host of *E. granulosus*, and 5.1-74.4% prevalence has been reported [11,12]. The sheep strain was shown to be the most common genotype of *E. granulosus sensu stricto*. The majority of camels are occasionally infected by the camel strain [1,13]. Both morphological and molec-

\*Received 1 January 2015, revised 16 November 2015, accepted 26 November 2015.

\*Corresponding author (Bmeshgi@ut.ac.ir)

© 2016, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ular approaches could together provide more accurate and reliable information about the nature and extent of variants within *E. granulosus* [14-16]. The objective of this study was to identify the morphological and molecular characteristics of *E. granulosus* derived from the wild sheep (*O. orientalis*) in Iran.

## MATERIALS AND METHODS

### Sample preparation

The hydatid cyst was collected from the liver of a dead wild sheep in Khojir National Park near Tehran (Fig. 1). Protoscolices were collected in sterile saline containing antibiotics (0.4 µg/ml penicillin and 100 µg gentamycin). The vitality of protoscolices was determined by their overall aspects (motility, activity, and presence of calcareous corpuscles) and negative methylene blue staining. A 5-month-old puppy which had received the required vaccinations and anthelmintics was infected orally with 6,000 protoscolices which were derived from the wild sheep. After 41 days, the dog was euthanized by an intravenous injection of an overdose of sodium pentobarbital. At the autopsy, adult *E. granulosus* worms were immediately collected from the small intestine, washed extensively in PBS (pH 7.4), and fixed in 70% ethanol for molecular evaluation.

### Morphological examinations

Protoscolices were mounted in polyvinyl lactophenol, and sufficient pressure was applied to the coverslip in order to flatten it without damaging the hooks. All measurements were taken on an Olympus microscope by the same individual using a calibrated ocular micrometer. The total length of large (LTL) and small (STL) hooks, blade length of large (LBL) and small (SBL) hooks, and the ratio of blade length to total length in large (LBL/LTL%) and small (SBL/STL%) hooks were measured. Three large and 3 small hooks were measured, and all hooks were counted from each of 6 protoscolices [17]. The ratio of blade length was the percentage of blade length to total length of large (LBL/LTL%) and small (SBL/STL%) hooks. The shape and arrangement of hooks were noticed.

### DNA extraction and PCR

Genomic DNA was extracted from an adult individual *E. granulosus* which was derived from wild sheep (*O. orientalis*) using DNA kit (MBST, Tehran, Iran) and following the manufacturer's recommendations. All the DNA samples were stored at -20°C until they were used. DNA fragments of ribosomal ITS2 and mitochondrial COX1 were amplified using the primers, including EgI: (forward; 5'-GTCTGTCCGAGCGTCGGCTTGIAA-3') and (reverse; 5'-CCCATCCACCACAGCATCCA-3') for ITS2, and




Fig. 1. Khojir National Park located in the northeast of Tehran, Iran.

EgC: (forward; 5'-TAGGGTTTTTATGACCAAAG-3') and (reverse; 5'-CACACCACATTTATCCAATT-3') for COX1. The PCR mixture was prepared in a total volume of 50 µl containing 25 µl of Taq master mix (Amplicon, Pullman, Washington, USA), 2 µl of forward primer and 2 µl of reverse primer (final concentration of 0.4 µM), 10 µl of DNA, and 11 µl of deionized distilled water in an automated thermocycler (Biorad Italia, Milano, Italy). The PCR was performed using the following protocol: 5 min incubation at 94°C to denature the double stranded DNA, 38 cycles for 45 sec at 94°C (denaturing step), staying for 45 sec at 44°C for COX1 primer or 45 sec at 56.6°C for ITS2 specific primers (annealing step), and 45 sec at 72°C (the extension step). Finally, the PCR was completed with an additional extension step for 10 min at 72°C. The DNA size marker (Vivantis, Romania) was used to estimate the length of the ITS2 or the COX1 amplicons. The PCR products were analyzed on 1.2% agarose gels in 0.5×TBE buffer and visualized using Sybersafe staining (Cinaclone, Tehran, Iran) and a UV illuminator. Consequently, the PCR product was purified using a quick PCR product purification kit (MBST) according to the manufacturer's recommendations.

### Sequencing and phylogeny

Sequencing was performed in both directions for each of the PCR products based on Sanger's method by the Kowsar Biotech Co. (Tehran, Iran), and was subjected to BLAST analysis through applying programs in the National Center for Biotechnology Information. The sequences were aligned and compared with each other and those of *E. granulosus* strains. The phylogenetic relationships of *E. granulosus* derived from

wild sheep isolates and based on ITS2 and COX1 were studied employing Mega 5.

Pairwise comparisons of the level of sequence differences (D) were conducted by the formula  $D = 1 - (M/L)$ , where M is the number of alignment positions at which the 2 sequences have a common base, and L is the total number of alignment positions with which the 2 sequences are compared [18].

### Statistical analysis

The results were statistically analyzed using the Student's *t*-test, and SPSS software (version 18) was utilized as well. Furthermore, the *P*-value of <0.05 was considered to be statistically significant.

## RESULTS

### Morphometric parameters

The hydatid cyst recovered from the liver of a wild sheep was fertile and had enough protoscolices for study on the morphology and genotype characters. The rostellar hooks of the protoscolices consisted of 2 rows with alternating regular arrangements of large and small hooks, and their outlines were rough. Comparison was carried out between the measurements of different variables from sheep-dog (*sensu stricto*), camel-dog (*intermedius*) from Iran [5], and wild sheep isolate. The results are summarized in Table 1.


The results showed that the total length and the blade length of large and small hooks are almost similar to those of *sensu stricto* strain, while they have statistically significant differences from those of the camel strain ( $P < 0.05$ ).

**Table 1.** Comparison between morphological characteristics of rostellar hooks of protoscolices of wild sheep, sheep, and camel isolates in Iran

| Characteristics of hooks | Wild sheep                        | Sheep-dog strain <sup>a</sup> | Camel-dog strain <sup>a</sup>     |
|--------------------------|-----------------------------------|-------------------------------|-----------------------------------|
| <b>Large hook</b>        |                                   |                               |                                   |
| Total length (LTL) (µm)  | 23.6±1.4 (22.4-25.2) <sup>b</sup> | 23.3±2.9 (21.4-24.8)          | 28.9±1.5 (24.8-32.5) <sup>b</sup> |
| Blade length (LBL) (µm)  | 10.7±1.1 (8.4-11.2) <sup>b</sup>  | 11.7±0.9 (10.1-13.5)          | 14.2±2.9 (12.4-31.5) <sup>b</sup> |
| LBL/LTL (%)              | 45.3±5.2 (33.0-50.0)              | 50.5±2.3 (42.1-54.5)          | 49.3±5.5 (42.3-58.3)              |
| <b>Small hook</b>        |                                   |                               |                                   |
| Total length (STL) (µm)  | 19.6±1.7 (16.7-22.4) <sup>b</sup> | 18.1±1.0 (16.9-20.3)          | 21.6±1.4 (18.0-23.0) <sup>b</sup> |
| Blade length (SBL) (µm)  | 6.6±1.3 (5.6-8.4) <sup>b</sup>    | 7.4±0.8 (6.7-9.0)             | 9.7±1.1 (6.7-11.0) <sup>b</sup>   |
| SBL/STL (%)              | 33.4±5.6 (28.5-43.0)              | 40.6±3.3 (33.3-45.2)          | 45.2±3.4 (30.0-48.0)              |
| Total number of hooks    | 34.8±1.9 (30.0-36.0)              | 35.5±4.2 (29.0-44.0)          | 33.7±2.6 (31.0-39.0)              |
| Outline                  | Rough                             | Rough                         | Smooth                            |
| Arrangement              | Alternate                         | Alternate                     | Alternate                         |

<sup>a</sup>Data from Hosseini and Eslami [5].

<sup>b</sup>Statistically significant differences ( $P < 0.05$ ) between wild sheep isolate of *E. granulosus* and camel strain.




**Fig. 2.** Agarose gel (1.2%), electrophoresis COX1, and ITS2 PCR products of *E. granulosus* derived from wild sheep isolates. Lane 1, ITS2 gene; lane 2, DNA size marker; lane 3, COX1 gene; lane 4, negative control.


**Molecular characteristics**

DNA was extracted from adult *E. granulosus* derived from the wild sheep. PCR was performed for 378 and 857 bp fragments of *E. granulosus* ITS2 and COX1 genes, respectively, using specific primers. The results are shown in Fig. 2. The distance based analysis, using neighbor-joining and parsimony analysis, was conducted for phylogenetic trees in order to evaluate the genetic relationships (Figs. 3, 4). The sequences were submitted to GenBank (accession no. KR872308 for ITS2 and KR905709 for COX1).

The similarity of these sequences with the recorded data in GenBank varied 99% for ITS2 and 98% for COX1 genes. The phylogenetic tree of *E. granulosus* from wild sheep isolate based on ITS2 was first computed altogether and then with an outgroup of *Dirofilaria immitis* to place the root (Fig. 3). The values on the nodes are bootstrap support from maximum likelihood and neighbor-joining analysis. The bootstrap supporting lower than 90% is not shown. COX1 nucleotide sequences were also used to construct a phylogenetic tree with an outgroup of *D. immitis* (Fig. 4). Two genotypes of *E. granulosus* corresponding to the camel and sheep isolates showed 10% differences, and wild sheep isolate was grouped into a



**Fig. 3.** Phylogenetic relationships of *E. granulosus* which was derived from wild sheep isolate in Iran based on ITS2 sequences in comparison with different isolates.



**Fig. 4.** Phylogenetic relationships of *E. granulosus* which was derived from wild sheep isolates in Iran based on COX1 sequences in comparison with different isolates.

distinct cluster corresponding to the sensu stricto genotype.

**DISCUSSION**

Parasites play an important role for regulation of host population in a natural environment. Therefore, having enough knowledge about helminthic infections of wildlife is essential in every area. Echinococcosis is endemic in Iran and is maintained in 3 distinct cycles; a livestock dog domestic cycle, a desert cycle between dogs and camels, and a sylvatic cycle between wild carnivores and wild ruminants [19]. The domestic cycle in Iranian sheep is the most important intermediate reservoir with the prevalence of 5.1-74.4% in sheep [11,12], 3.3-

63.3% in farm dogs, and 5.0-50.5% in stray dogs [11,20]. In the sylvatic cycle, 5.0-8.7% of golden jackals [3,4] act as the final host. Intermediate hosts, including wild sheep and wild pigs [2], were found to be infected with *E. granulosus* in this cycle. The stray dogs, farm dogs, and wild carnivores are in close contact with farmers, their families, and their residential areas as well as the nomads and their black trends mainly during their wandering. Free-roaming dogs and other carnivores may be fed on or scavenge for discarded offal of domestic and wild ruminants like camels.

Drawing the hooks of protoscolices for measurements has several advantages. The total and handle lengths of both large and small hooks are considered as the most variable characteristics that could be used not only for differentiating parasite isolates from different host species but also for the original infection in the definitive host. Due to intraspecific variation of sheep and camel strains, using both morphologic and molecular tools concomitantly provides more accurate and reliable information about the phenotype and genotype characteristics of *E. granulosus* isolates and emphasizes the valuable tools in transmission and epidemiological studies as well.

In this study, the liver of a wild sheep infected with hydatid cysts from the Khojir National Park near Tehran was used in determining the *E. granulosus* phenotype. It should be noted that this wild sheep isolate was highly similar to the sensu stricto strain ( $P < 0.05$ ), which is highly important from the zoonotic point of view. The possibility of interaction between the pastoral and sylvatic cycles of *E. granulosus* [14-16] which were found in Khojir National Park, Tehran, Iran in an examined wild sheep suggested the possibility of interaction of 3 different cycles existing worldwide.

The genotyping of 16 isolates of *E. granulosus* from humans, sheep, goats, cattle, and camels revealed the existence of 2 distinct strains (sheep-dog and camel-dog strains) by mtDNA markers [9]. As molecular analyses of the sylvatic cycle of echinococcosis, NADH dehydrogenase subunit 1 (ND1) and COX1 sequences were used to characterize deer isolates of *E. granulosus* from Canada and Finland, respectively [21,22]. In the second part of the present study, ITS2 and COX1 genes of *E. granulosus* derived from an Iranian wild sheep were amplified and sequenced. Our findings showed the sequence of fragments of 378 and 857 bp for ITS2 and COX1 genes, respectively. Two nucleotide differences of 5.1% (16/314) and 5.4% (16/295) were obtained for ITS2 and COX1 which were at 9 and 8 variable sequence positions, respectively. The COX1

nucleotide sequence obtained from deer isolate [22] was different from what our findings (92% similarity) demonstrated. ITS2 and COX1 sequences indicated that *E. granulosus* derived from the present wild sheep could be categorized as the sheep-dog genotype. Moreover, no variation was observed in the composition of the 2 obtained gene sequences (KR872308 and KR905709) deposited in GenBank.

In conclusion, our results showed that both morphological characters and molecular analyses were confirmed to be complementary for determination of *E. granulosus* which was derived from an Iranian wild sheep (*O. orientalis*) as a sensu stricto strain in Iran. To sum up, treatment of non-human intermediate hosts is not practical, although the frequented baiting in Australian wild definitive hosts containing praziquantel is likely to be practical [23]. Human beings can ingest eggs if they fail to follow sanitary and hygienic practices, including wearing protective barriers like gloves and washing their hands well with soap and water after possible exposure to canid feces. In humans, this exposure could be most effectively curtailed by preventing the consumption of cervid viscera by free-roaming dogs. The parasites such as *E. granulosus* strains should be identified and monitored. The results of the present study showed that the sensu stricto strain is also an infective strain in the wildlife, and the wildlife infection reflects the infections of domestic animals.

## ACKNOWLEDGMENTS

Project supports were provided by the University of Tehran, Iran (grant no. 3/1/7502004). The authors are very grateful to Environmental Organization for help.

## CONFLICT OF INTEREST

We have no conflict of interest related to this work.

## REFERENCES

1. Fasihi Harandi M, Hobbs RP, Adams PJ, Mobedi I, Morgan-Ryan UM, Thompson RCA. Molecular and morphological characterization of *Echinococcus granulosus* of human and animal origin in Iran. *Parasitology* 2002; 125: 367-373.
2. Eslami A, Farsad-Hamdi S. Helminth parasites of wild boar, *Sus scrofa*, in Iran. *J Wildl Dis* 1992; 28: 316-318.
3. Meshgi B, Eslami A, Bahonar AR, Kharazian-Moghadam M, Gerami-Sadeghian A. Prevalence of parasitic infections in the

- red fox (*Vulpes vulpes*) and golden jackal (*Canis aureus*) in Iran. *Iran J Vet Res Shiraz Univ* 2002; 10: 387-391.
4. Sadighian A. Helminth parasites of stray dogs and jackals in Shahsavari area, Caspian Sea region. *J Parasitol* 1969; 55: 372-374.
  5. Hosseini SH, Eslami A. Morphological and developmental characteristics of *Echinococcus granulosus* derived of sheep, cattle and camel in Iran. *J Helminthol* 1998; 72: 337-341.
  6. Thompson RCA, McManus DP. Towards a taxonomic revision of the genus *Echinococcus*. *Trend Parasitol* 2002; 18: 452-457.
  7. Zhang L, Eslami A, Hosseini SH, McManus DP. Indication of the presence of two distinct strains of *Echinococcus granulosus* in Iran by mitochondrial DNA markers. *Am J Trop Med Hyg* 1998; 59: 171-174.
  8. Nakao M, McManus DP, Schantz PM, Craig PS, Ito A. A molecular phylogeny of the genus *Echinococcus* inferred from complete mitochondrial genomes. *Parasitology* 2007; 134: 713-724.
  9. Moks E, Jogisalu I, Valdmann H, Saarma U. First report of *Echinococcus granulosus* G8 in Eurasia and a reappraisal of the phylogenetic relationships of 'genotypes' G5-G10. *Parasitology* 2008; 135: 647-654.
  10. Aminpour A, Hosseini SH, Shayan P. Comparative genotyping of *Echinococcus granulosus* infecting buffalo in Iran using *cox1* gene. *Parasitol Res* 2011; 108: 1229-1234.
  11. Sharifi I. Seasonal prevalence of hydatid cyst in slaughterhouse of the city of Kerman. *Iran J Public Health* 1996; 25: 39-46.
  12. Daryani A, Alaei R, Arab R, Sharif M, Dehghan MH, Ziaei H. The prevalence, intensity and viability of hydatid cyst in slaughtered animals in the Ardabil province of Iran. *J Helminthol* 2007; 81: 13-17.
  13. Sharbatkhori M, Mirhendi H, Fasihi Harandi M, Rezaeian M, Mohebbi M, Eshraghian M, Rahimi H, Bigomkia E. *Echinococcus granulosus* genotypes in livestock of Iran indicating high frequency of G1 genotype in camel. *Experiment Parasitol* 2010; 124: 373-379.
  14. Eckert J, Thompson, RCA. Intraspecific variation of *Echinococcus granulosus* and related species with emphasis on their infectivity to humans. *Acta Trop* 1997; 64: 19-34.
  15. McManus DP, Bryant C. Biochemistry, physiology and molecular biology of *Echinococcus*. In Thompson RCA, Lymbery AJ eds, *Echinococcus and Hydatid Disease*. Walingford, Oxon, UK. CAB International. 1995, pp. 135-181.
  16. McManus DP, Bowles J. Molecular genetic approaches to parasite identification: their value in diagnostic parasitology and systematics. *Int J Parasitol* 1996; 26: 687-704.
  17. Hobbs RP, Lymbery AJ, Thompson RCA. Rostellar hook morphology of *Echinococcus granulosus* (Batsch, 1786) from natural and experimental Australian hosts and its implications for strain recognition. *Parasitology* 1990; 101: 273-281.
  18. Chilton NB, Gasser RB, Beveridge I. Differences in a ribosomal DNA sequence of morphologically indistinguishable species within the *Hypodontus macropi* complex (Nematoda: Strongyloidea). *Int J Parasitol* 1995; 25: 647-651.
  19. Ahmadi N, Dalimi A. Characterization of *Echinococcus granulosus* isolates from human, sheep and camel in Iran. *Infect Genet Evol* 2006; 6: 85-90.
  20. Eslami A, Hoseini SH. *Echinococcus granulosus* infection of farm dogs of Iran. *Parasitol Res* 1998; 84: 205-207.
  21. Himsforth CG, Jenkins E, Hill JE, Nsungu M, Ndao M, Thompson RCA, Covacin C, Ash A, Wagner BA, McConnel, A, Leighton FA, Skinner S. Emergence of sylvatic *Echinococcus granulosus* as a parasitic zoonosis of public health concern in an indigenous community in Canada. *Am J Trop Med Hyg* 2010; 82: 643-645.
  22. Lavikainen A, Lehtinen MJ, Laaksonen S, Agren E, Oksanen A, Meri S. Molecular characterization of *Echinococcus* isolates of cervid origin from Finland and Sweden. *Parasitology* 2006; 133: 567-570.
  23. Claridge AW, Mills DJ, Hunt R, Jenkins DJ, Bean J. Satellite tracking of wild dogs in south-eastern mainland Australian Forests: implications for management of a problematic top-order carnivore. *Forset Ecol Manag* 2009; 256: 814-822.