



Subsets of Inflammatory Cytokine Gene Polymorphisms are Associated with Risk of Carcinogenic Liver Fluke *Opisthorchis viverrini*-Associated Advanced Periductal Fibrosis and Cholangiocarcinoma

Arpa Surapaitoon^{1,2}, Sutas Suttiprapa^{2,3}, Eimorn Mairiang⁴, Narong Khuntikeo⁵, Chawalit Pairojkul⁵, Jeffrey Bethony⁶, Paul J. Brindley⁶, Banchob Sripa^{2,3,*}

¹Biomedical Sciences, Graduate School, Khon Kaen University, Khon Kaen, Thailand; ²WHO Collaborating Centre for Research and Control of Opisthorchiasis (Southeast Asian Liver Fluke Disease), Tropical Disease Research Center; ³Department of Pathology; ⁴Department of Radiology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand; ⁵Liver Fluke and Cholangiocarcinoma Research Center, and Department of Surgery, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand; ⁶Department of Microbiology, Immunology and Tropical Medicine, and Research Center for Neglected Tropical Diseases of Poverty, School of Medicine & Health Sciences, The George Washington University, Washington DC, 20037, USA

Abstract: *Opisthorchis viverrini* infection induces chronic inflammation, and a minor proportion of infected individuals develop advanced periductal fibrosis (APF) and cholangiocarcinoma (CCA). Inflammatory cytokines and/or their gene polymorphisms may link to these biliary pathologies. We therefore investigated associations among cytokine gene polymorphisms and cytokine production in 510 Thai cases infected with *O. viverrini* who presented with APF+ or APF-, as established by abdominal ultrasonography as well as in patients diagnosed with CCA. Levels of pro-inflammatory and anti-inflammatory cytokines were determined in culture supernatants after stimulation of peripheral blood mononuclear cells (PBMCs) with *O. viverrini* excretory-secretory (ES) products. Pro-inflammatory cytokines, IL-1 β , IL-6, IFN- γ , LT- α , and TNF- α were significantly increased in CCA patients compared with non-CCA (APF- and APF+) cases. Polymorphisms in genes encoding IL-1 β -511C/T, IL-6 -174G/C, IFN- γ +874T/A, LT- α +252A/G, and TNF- α -308G/A were then investigated by using PCR-RFLP or allele specific-PCR (AS-PCR) analyses. In the CCA cases, LT- α +252A/G and TNF- α -308G/A heterozygous and homozygous variants showed significantly higher levels of these cytokines than the wild type. By contrast, levels of cytokines in wild type of IFN- γ +874T/A were significantly higher than the variants in CCA cases. IFN- γ +874T/A polymorphisms were associated with advanced periductal fibrosis, whereas IL-6 -174G/C polymorphisms were associated with CCA. To our knowledge, these findings provide the first demonstration that *O. viverrini* infected individuals carrying several specific cytokine gene polymorphisms are susceptible to develop fibrosis and CCA.

Key words: *Opisthorchis viverrini*, hepatobiliary disease, advanced periductal fibrosis, cholangiocarcinoma, cytokine gene polymorphisms

INTRODUCTION

Chronic infection with the carcinogenic liver fluke *Opisthorchis viverrini* results in continuous irritation, chronic inflammation, restlessness tissue damage, and repair of the bile tract, including the second order bile ducts [1]. Consequently, infection with *O. viverrini* is strongly associated with chronic hepatobiliary disease, including cholangitis, cholecystitis, obstruc-

tive jaundice, advanced periductal fibrosis, and cholangiocarcinoma (CCA) [1,2]. Ultrasonographic assessment of hepatobiliary abnormalities of otherwise asymptomatic persons stool positive for infection with *O. viverrini* revealed that about 25% of these cases developed advanced periductal fibrosis, and CCA was suspected in up to 0.5% of them based on abdominal ultrasound examinations [3,4]. The mechanism underlying the different outcomes of *O. viverrini* infection is unknown. Our extensive research experience with opisthorchiasis led us to suspect that host inflammatory responses to the liver fluke might play pivotal roles in the outcome [1,2]. Inflammation mediated by cytokines has been described in infectious diseases and malignancies [5,6]. Specifically, the liver fluke infection induces marked inflammation characterized by infiltration by

•Received 18 April 2017, revised 8 May 2017, accepted 11 May 2017.

*Corresponding author (banchob@kku.ac.th)

© 2017, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

neutrophils and mononuclear cells into the vicinity of the infected bile ducts. Proteins and other metabolites released by the *O. viverrini* worms activate Toll-like receptor 4 and induce expression of the potent chemo-attractant IL-8 and pro-inflammatory IL-6 cytokines [7].

Gene polymorphisms affect the cytokine expression and influence inflammatory processes in response to infectious diseases and cancer [8]. In the context of liver fluke infection, hepatobiliary disease and associated bile duct cancer, we have reported marked elevation in expression of IL-6 in response to *O. viverrini* excretory-secretory (ES) antigens during chronic opisthorchiasis, particularly where biliary periductal fibrosis is marked [9]. In addition, we observed significant increases in both pro-inflammatory and anti-inflammatory cytokines during cholangiocarcinoma associated with chronic liver fluke infection [10]. We now hypothesize that cytokine gene polymorphisms influence cytokine expression profiles, which in turn may result in diverse pathogenic sequelae during infection with *O. viverrini*. This study described below investigated associations among cytokine gene polymorphisms and cytokine production during chronic opisthorchiasis and associated biliary tract diseases, including CCA.

MATERIALS AND METHODS

Study participants

This study was approved by the Khon Kaen University Human Ethics Committee, reference nos. HE480528 and HE 581272. Written informed consents were obtained from all participants or patients. Four hundred participants positive for infection with *O. viverrini*, including 200 without APF- and 200 with APF+, were recruited from sites endemic for *O. viverrini* infection in Khon Kaen or neighboring provinces in north-eastern Thailand, as described [9]. This study also included 110 histologically proven cases of cholangiocarcinoma (CCA); these latter participants had undergone hepatectomy at Srinagarind Hospital, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand.

Opisthorchis viverrini-ES product preparation

ES products of *O. viverrini* (Ov-ES) were prepared as described [11,12]. Briefly, viable liver flukes recovered at necropsy from experimentally infected hamsters were cultured in RPMI-1640 containing antibiotics and the protease inhibitors (0.1 mM phenylmethanesulfonyl fluoride (PMSF), 1 mM leu-

peptin and 0.1 mM N-[N-(L-3-trans-carboxyoxiran-2-carboxyl)-L-leucine]-agmatine, E-64). The cultures were maintained in vitro at 37°C and supernatants containing the Ov-ES were collected twice daily for 7 days. The culture supernatants were subjected to centrifugation at 3,000 g for 10 min to pellet the eggs. Clarified supernatants were pooled, dialyzed against PBS, and concentrated by membrane ultrafiltration (PM 10, Amicon, Danvers, Massachusetts, USA). Lipopolysaccharide (LPS) was removed using Triton-X114 extraction [13], followed by passage through Bio-Beads SM2 (Bio-Rad, Hercules, CA, USA) to remove Triton-X114. Finally, OvES was filtered through a 0.2 µm membrane and then aliquoted and stored at -80°C. The LPS concentration was determined using the *Limulus* amoebocyte assay (less than 100 ng/ml failed to affect cell proliferation or cytokine production, unpublished) [14].

Peripheral blood mononuclear cells (PBMCs) culture

PBMC were isolated from heparinized blood by density gradient centrifugation using Ficoll-Paque (Sigma Diagnostic, St. Louis, Missouri, USA), counted on a hemacytometer and viability assessed using 0.5% Trypan blue dye. PBMC were cultured at 2×10^4 cells/ml in RPMI-1640 supplemented with antibiotics and 5% fetal bovine serum with or without Ov-ES, 20 µg/ml. Phytohemagglutinin (PHA)-L (Difco Laboratories, Detroit, Michigan, USA) at 10 µg/ml was used as positive control of cytokine stimulation. The cells were cultured at 37°C in a humidified 5% CO₂ incubator for 48 hr. Supernatants were harvested and stored at -80°C for cytokine level determination.

Cytokine detection

A 25-50 µl of supernatant was used for cytokine assays. The secreted cytokines were quantified using the FlowCytomix bead-based multiplexing assays kit, human Th1/Th2 (IFN-γ, IL-1β, IL-2, IL-4, IL-5, IL-6, IL-8, IL-10, IL-12p70, TNF-α, and LT-α) 11-plex kit (BMS810FF, BenderMedSystems, Vienna, Austria) and quantified using flow cytometry (FC500, Beckman-Coulter, Brea, California, USA). Cytokine levels were established by interpolation from standard curves established using human recombinant cytokines (BMS8400FF, BenderMedSystems), after which the findings were expressed as pg/ml.

DNA extraction and genotyping

Genomic DNA was extracted from PBMC from each participant using the Puregene DNA Isolation Kit (Gentra systems, Minneapolis, Minnesota, USA), quantified using the Nano-

Table 1. Primer sequences, restriction enzymes, and size of digestion products in the PCR-RFLP and AS-PCR analyses

Gene/SNP	rsn	wt/var	Region	Oligonucleotide primers	RE	Digestion products (bp)
IL-1β/-511	rs16944	C/T	Promoter	F (5'- TGG CAT TGA TCT GGT TCA TC-3') R (5'-GTT TAG GAA TCT TCC CAC TT-3')	Ava I	wt: 190, 114 var: 304
IL-6/-174	rs1800795	G/C	Promoter	F (5'-GGG CTG CGA TGG AGT CGA AG-3') R (5'-TCC CTC ACA CAG GGC TCG-3')	Nla III	wt: 233, 54 var: 122, 111, 54
IFN-γ/+874	rs2430561	T/A	Intron 1	Common primer: 5'TCAACAAAGCTGATACTCCA-3', T allele primer: 5'TTCTTACAACACAAAATCAAATCT-3', A allele primer: 5'TCTTACAACACAAAATCAAATCA-3'		
LT-α/+252	rs909253	A/G	intron	F (5'-CCG TGC TTC GTG CTT TGG ACT A-3') R (5'-AGA CGT TCA GGT GGT GTA AT-3')	Nco I	wt: 427 var: 231, 196
TNF-α/-308	rs1800629	G/A	promoter	F (5'-TCC TCC CTG CTC CGA TTA CG-3') R (5'-AGG CAA TAG GTT TTG AGG GCC-3')	Nco I	wt: 87 var: 107

SNP, Single nucleotide polymorphism; rsn, reference SNP number; wt, wild type; var, variant; RE, restriction enzyme; bp, base pair




Fig. 1. Representative amplicons and digestion fragments corresponding to gene polymorphisms. (A) 190 and 114 bp fragments specific for C allele, 304 bp fragment specific for T allele of IL-1b-511C/T polymorphism. (B) 233 bp and 54 bp bands specific for G allele, 122 bp, 111 bp, and 54 bp bands specific for C allele of IL-6-174G/C polymorphism. (C) 262 bp fragment specific for T and A alleles of IFN-γ +874T/A polymorphism. (D) 427 bp fragment specific for A allele, 231 and 196 bp fragments specific for G allele of LT-α +252A/G polymorphism. (E) 107 bp fragment specific for A allele, 87 bp fragments specific for G allele of TNF-α -308G/A polymorphism. M, 50 bp DNA marker.

Drop ND-1000 and stored at -20°C. Single nucleotide polymorphisms (SNPs) for IL-1β -511C/T, IL-6 -174G/C, LT-α

+252A/G, and TNF-α -308G/A polymorphisms were analyzed by restriction fragment length polymorphism (RFLP), as de-

scribed [15-18]. Allele specific-PCR (AS-PCR) was used to analyze for IFN- γ +874T/A polymorphism [19]. Table 1 lists the PCR thermal cycling conditions and oligonucleotide primers specific for 5 cytokine gene polymorphisms, IL-1 β -511C/T, IL-6 -174G/C, IFN- γ +874T/A, LT- α +252A/G, and TNF- α -308G/A. Amplified products were evaluated by 1.5% (IFN- γ), 3.0% (IL-6, LT- α and TNF- α), and 8.0% (IL-1 β) agarose gel electrophoresis containing ethidium bromide (0.5 μ g/ml). After electrophoresis, the ethidium-stained fragments were visualized and documented (Multi Doc-It™ Digital Imaging System, UVP, Upland, California, USA) (Fig. 1).

Statistical analysis

The Kolmogorov-Smirnov test was used to test whether the data were distributed normally. Statistical analysis was performed with non-parametric tests (Kruskal-Wallis and Mann-Whitney), with assistance of the IBM SPSS Statistics for Windows package, version 19 (SPSS, Chicago, Illinois, USA). Hardy-Weinberg equilibrium (HWE) analyses (chi-square tests, χ^2) were used to compare the expected and observed genotype frequencies in the non-CCA groups of study participants. Allele frequency differences between cases (CCA) and controls (non-CCA) were analyzed, and the odd ratios (OR) and 95% confidence intervals (95% CI) were calculated.

RESULTS

Demographic and clinicopathological characteristics

Table 2 summarizes the clinicopathological characteristics of participants in this study who had been diagnosed with CCA as well as the liver fluke infected cases without CCA. The APF- group included 109 females and 91 males, mean age of 46.8 years (range, 20-60 years) and the APF+ group included 108 females and 92 males, mean age of 47.4 years (range, 25-60 years). The CCA group was composed of 72 men and 38 women, and the mean age was 55.9 years (range, 40-76 years).

Cytokine production in *Ov*-ES stimulated PBMCs

Fig. 2 summarizes the net production of all cytokines quantified following *Ov*-ES stimulation of PBMC from the APF-, APF+, and CCA cases. Levels of IL-1 β , IFN- γ , and LT- α were significantly higher in CCA patients compared with the APF- and APF+ groups. TNF- α production was significantly higher in CCA and APF+ groups compared with APF-. Notably, IL-6 was significantly higher in APF+ cases compared with the CCA

Table 2. Clinicopathological characteristics of cholangiocarcinoma (CCA) patients and non-CCA controls

Demographics	APF- (n=200)	APF+ (n=200)	CCA (n=110)
Sex			
Male (%)	91 (45.5)	92 (46.0)	72 (65.5)
Female (%)	109 (54.5)	108 (54.0)	38 (34.5)
Age (years)			
Mean \pm SD	46.8 \pm 7.7	47.4 \pm 7.5	55.9 \pm 10.3
Range	20-60	22-60	40-76
Age (in year, %)			
\leq 30	6 (3.0)	5 (2.5)	1 (0.9)
31-40	34 (17.0)	33 (16.5)	5 (4.6)
41-50	86 (43.0)	83 (41.5)	20 (18.2)
\geq 51	74 (37.0)	79 (39.5)	84 (46.4)

and APF- groups. Differences were not apparent among the groups for 6 other cytokines, IL-2, IL-4, IL-5, IL-8, IL-10, and IL-12p70.

Gene polymorphism and cytokine production

Overall comparison regardless of CCA or non-CCA groups did not reveal significant differences in levels of cytokines among gene polymorphisms except for LT- α , where net cytokine production in LT- α +252A/G heterozygous (AG) and homozygous (GG) variants were significantly elevated compared to the wild type (Fig. 3). However, when subdividing into CCA and non-CCA groups (APF- and APF+), LT- α +254A/G heterozygous (AG), and homozygous (GG) variants showed significantly higher levels of cytokines than the wild type in CCA group; 2.35-fold and 2.24-fold, respectively. Similarly, TNF- α -308G/A heterozygous (GA) and homozygous (AA) variants showed significantly higher levels of cytokines than the wild type in CCA group with fold changes of 12.49 and 4.76, respectively. Interestingly, IFN- γ +874T>A heterozygous (TA) and homozygous (AA) was significantly lower (0.56- and 0.2-fold, respectively) than the wild type in the CCA group.

Distribution of cytokine gene polymorphisms among patient groups

Table 4 presents the genotypes and allele distribution of IL-1 β -511C/T, IL-6 -174G/C, IFN- γ +874T/A, LT- α +254A/G, and TNF- α -308G/A polymorphisms in the participants, those with APF-, APF+, and CCA. Comparison among groups did not reveal significant differences in the genotype or allele frequencies for IL-1 β -511C/T, LT- α +252A/G, and TNF- α -308G/A. However, comparing the IL6 -174G/C genotypes between the CCA and non-CCA (APF- or APF+) groups revealed that the presence of heterozygous (GC) genotype associated with CCA




Fig. 2. Net cytokine profile of OvES stimulated PBMCs in APF-, APF+, and CCA patients. The Wilcoxon Matched-paired Signed rank test was used to assess the statistical significance; * $P < 0.05$.

(OR=2.35 and 2.95 for APF- vs CCA and APF+ vs CCA, respectively). Moreover, a stronger association with CCA was found with the homozygous variant (CC) (OR= 5.50 and 7.52 for APF- vs CCA and APF+ vs CCA, respectively). In addition, C alleles at this locus also associated with CCA (OR=3.36) when the CCA and APF+ groups were compared. Comparing between APF+ and APF- groups, IFN- γ +874T>A heterozygous (TA) and homozygous (AA) variants associated with biliary periductal fibrosis (OR= 3.10 and 2.74, respectively). The IFN- γ +874T>A homozygous variant (AA) genotype was also associated with CCA (OR=2.20) as compared between APF- and CCA groups, whereas the A allele associated with both fibrosis (OR= 1.60, APF- vs APF+) and CCA (OR= 1.71, APF- vs CCA). In the case of TNF- α -308G/A polymorphism, only the TNF- α -308 A allele appeared to be associated with CCA (OR= 1.61 and OR= 1.5 in APF- vs CCA and in APF+ vs CCA, respectively).

Table 3. Cytokine levels as established in PBMCs pulsed with excretory secretory products of the liver fluke *Opisthorchis viverrini* and the net cytokine production and genotype polymorphisms

	Cytokine levels, pg/ml		
	Wild type	Heterozygous	Variant
IL-1 β	268.4 \pm 47.7	315.4 \pm 35.9	303.1 \pm 41.1
IL-6	3,973.1 \pm 763.2	4,221.6 \pm 434.8	4,323.7 \pm 440.5
IFN- γ	514.9 \pm 140.3	397.5 \pm 117.4	239.9 \pm 65.2
LT- α	67.3 \pm 17.9	187.4 \pm 36.0*	153.7 \pm 36.7*
TNF- α	362.6 \pm 68.8	498.8 \pm 127.1	466.7 \pm 67.9

Cytokine levels presented as mean \pm SD; * $P < 0.05$; n=510.

DISCUSSION

Infection with the liver fluke *O. viverrini* frequently leads to chronic inflammation of the bile ducts and results in unremitting tissue damage that, in turn, can cause hepatobiliary dis-




Fig. 3. The net production of IL-1β, IL-6, IFN-γ, LT-α, and TNF-α, and their genotype polymorphisms in APF- (n=200), APF+ (n=200), and CCA patients (n=110). wt, wild type; hetero, heterozygous; var, variant.

Table 4. Comparison of genotypes and allele frequencies for IL-1 β , IL-6, IFN- γ , LT- α , and TNF- α in CCA cases and the participants in the APF+ and the APF- groups

Cytokine/ Genotype/Allele	APF-N(%)	APF+N(%)	CCA N(%)	APF- vs. APF+	APF- vs. CCA	APF+ vs. CCA
				OR (95% CI)	OR (95% CI)	OR (95% CI)
IL-1β (-511C/T)						
CC	38 (19.0)	36 (18.0)	18 (16.1)	1	1	1
CT	100 (50.0)	93 (46.5)	48 (43.6)	0.98 (0.50- 1.74)	1.01 (0.50-2.09)	1.03 (0.50-2.14)
TT	62 (31.0)	71 (35.5)	44 (40.0)	1.20 (0.65-2.22)	1.49 (0.72-3.16)	1.23 (0.59-2.61)
C	176 (44.0)	165 (41.2)	84 (38.2)	1	1	1
T	244 (56.0)	235 (58.8)	136 (61.8)	1.02 (0.77-1.36)	1.16 (0.82-1.65)	1.13 (0.80-1.61)
IL-6 (-174G/C)						
GG	112 (56.0)	124 (62.0)	32 (29.1)	1	1	1
GC	67 (33.5)	59 (29.5)	45 (40.9)	0.79 (0.50-1.25)	2.35 (1.31-4.21)*	2.95 (1.64-5.31)*
CC	21 (10.5)	17 (8.5)	33 (30.0)	0.73 (0.34-1.53)	5.50 (2.65-11.41)*	7.52 (3.52-16.19)*
G	291 (72.7)	307 (76.7)	109 (49.5)	1	1	1
C	109 (27.3)	93 (23.3)	111 (50.5)	0.81(0.58-1.13)	0.26 (0.12-0.52)	3.36 (2.32-4.85)*
IFN-γ (+874T/A)						
TT	61 (30.5)	26 (13.0)	23 (20.9)	1	1	1
TA	74 (37.0)	98 (49.0)	33 (30.0)	3.10 (1.73-5.61)*	1.18 (0.60-2.34)	0.38 (0.18-0.80)
AA	65 (32.5)	76 (38.0)	54 (49.1)	2.74 (1.50-5.04)*	2.20 (1.13-4.20)*	0.80 (0.39-1.64)
T	196 (49.0)	150 (37.5)	79 (35.9)	1	1	1
A	204 (51.0)	250 (62.5)	141 (64.1)	1.60 (1.19-2.14)*	1.71 (1.20-2.44)*	1.07 (0.75-1.5)
LT-α (+252A/G)						
AA	60 (30.0)	47 (23.5)	24 (21.8)	1	1	1
AG	74 (37.0)	93 (46.5)	54 (49.1)	1.60 (0.95-2.65)	1.84 (0.97-3.45)	1.17 (0.60-2.16)
GG	66 (33.0)	60 (30.0)	32 (29.1)	1.16 (0.66-2.01)	1.21 (0.61-2.40)	1.04 (0.51-2.11)
A	194 (48.5)	187 (46.8)	102 (46.4)	1	1	1
G	206 (51.5)	213 (53.2)	118 (53.6)	1.07 (0.80-1.43)	1.08 (0.77-1.53)	1.01 (0.72-1.43)
TNF-α (-308G/A)						
GG	34 (17.0)	35 (17.5)	17 (15.5)	1	1	1
GA	54 (27.0)	46 (23.0)	13 (11.8)	0.82 (0.42-1.60)	0.48 (0.18-1.20)	0.58 (0.22-1.47)
AA	112 (56.0)	119 (59.5)	80 (72.7)	1.03 (0.58-1.83)	1.42 (0.71-2.92)	1.38 (0.69-2.82)
G	122 (30.5)	116 (29.0)	47 (21.4)	1	1	1
A	278 (69.5)	284 (71.0)	173 (78.6)	1.07 (0.78-1.47)	1.61 (1.08-2.43)*	1.50 (1.00-2.26)*

*Significant at *P*-value<0.05.

eases including CCA. A subset of people with opisthorchiasis develop overt hepatobiliary disease, with about 25% of them displaying marked biliary tract fibrosis and as many as 0.5% of infected cases exhibiting incipient bile duct cancer upon ultrasound examination [4,20,21]. Why infection leads to different disease outcomes among the population in high liver fluke infection endemic sites remains to be established. Divergent immunological responses among infected persons may contribute to the outcome, in addition to other co-factors [22]. The pro-inflammatory cytokine IL-6 is significantly elevated in infected individuals with marked and chronic hepatobiliary fibrosis [9,23] and cholangiocytes upregulate expression of IL-6 and IL-8 in response to the *Ov*-ES components [24,25]. Moreover, the recent study revealed both pro-inflammatory and anti-inflammatory cytokines were increased in the *O. viverrini*-associated CCA compared to uninfected normal controls [10].

Here, we investigated pro-inflammatory and anti-inflamma-

tory cytokine profiles among different groups of patients infected with *O. viverrini*: liver fluke infected individuals exhibiting biliary tract disease (advanced periductal fibrosis, APF+) or without overt biliary periductal fibrosis (APF-), and liver fluke infection-associated CCA cases. We observed that pro-inflammatory cytokines, IL-1 β , IL-6, IFN- γ , LT- α , and TNF- α , though no other cytokines were increased significantly during CCA in comparison with participants infected with *O. viverrini* but without CCA, i.e., the APF- and APF+ groups. These findings indicated that marked inflammatory responses during opisthorchiasis increased the risk for cholangiocarcinogenesis.

Gene polymorphisms are known to affect cytokine levels [8,26,27]. Single nucleotide polymorphism (SNP) can affect gene function in many ways [28]. If it occurs at the non-coding region, it can affect the binding of transcriptional factors, interfere RNA splicing, and increase or decrease messenger RNA degradation. SNP in coding region can result in 2 consequenc-

es; synonymous and nonsynonymous SNPs. Synonymous SNPs do not affect the protein sequence, but can still affect the protein conformation such as the silent polymorphism in the MDR1 gene changes substrate specificity [29]. Nonsynonymous SNPs change the amino acid sequence of protein and subsequent protein function as shown in cystic fibrosis patients [30]. For SNPs in our liver fluke associated CCA, we report here that certain cytokine gene polymorphisms affect proinflammatory cytokine production. However, detail molecular mechanisms need to be further investigated.

Accordingly, we hypothesized that cytokine gene polymorphisms may underline variations in inflammatory cytokine response in response to liver fluke infection and downstream biliary tract inflammation and cholangiocarcinogenesis. The gene polymorphisms IL-1 β -511C/T, IL6 -174G/C, LT- α +252A/G, and TNF- α -308G/A generated the high-producer phenotypes, whereas IFN- γ +874T/A exhibited a low-production phenotype. In agreement with these reports, both LT- α +252A/G and TNF- α -308G/A heterozygous and homozygous variants showed significant higher levels of cytokine than the wild type. Similarly, both IL-1 β -511C/T and IL-6 -174G/C variants produced more cytokines than the wild type, even though the difference was not statistically significant. By contrast, the variants of IFN- γ +874T/A produced less cytokines than the wild type. We note, however, that these cytokine production profiles were observed only in the CCA cases, and not among the non-CCA groups. The difference of cytokine production from the same gene polymorphism between non-CCA and CCA patients might indicate divergence in cytokine regulation/networking among the groups in this study.

Subsequently we investigated whether gene polymorphisms of these pro-inflammatory cytokine were related to the development of fibrosis or CCA. An IL-6 -174G>C polymorphism represents a potential modulator of risk for malignancy with ethnicity and tumor site dependent trends [31]. The present findings revealed that both heterozygous (GC) and homozygous (CC) variant of IL-6 -174G>C polymorphisms were associated with cholangiocarcinogenesis in this Thai population. Given that the heterozygous (GC) and homozygous (CC) variants of IL-6 -174G>C are high producer phenotypes, these gene polymorphisms might explain elevated levels of IL-6 during in CCA, as reported earlier. The A allele of the TNF- α -308G/A polymorphism appears to associate with the disease progression of hepatitis B viral infection and hepatocellular carcinoma [32]. In likely fashion, the present findings revealed

that the A allele of TNF- α -308G/A polymorphism associated with CCA.

Interferon gamma plays key roles in the immune responses and in antitumor mechanisms [33,34]. IFN- γ +874T>A polymorphisms are associated with viral and bacterial infections, which result in chronic inflammation and carcinogenic [35-37]. Here, IFN- γ +874T>A heterozygous (TA) and homozygous (AA) variants associated with fibrosis, and the IFN- γ +874T>A homozygous (AA) variant was also associated with CCA. As noted, these variants produce less IFN- γ than the TT (wild type) genotype. Together these findings indicate that in a subset of liver fluke infected individuals carrying these alleles, the phenotype of which is reduced levels of IFN- γ , are at elevated risk for biliary periductal fibrosis and CCA. Mechanisms underlying the association are not known. However, for anti-tumor activity, IFN- γ plays a central role in the recognition and elimination of transformed cells through its cytostatic, proapoptotic, and immune-provoking effects [33].

To summarize, cytokine and gene polymorphism profiles of Thais infected with the liver fluke *O. viverrini* were characterized. To our knowledge, this is the first demonstration that persons infected with *O. viverrini* inherited with certain gene polymorphisms that result in aberrant cytokine production are susceptible to develop fibrosis and CCA. Pro-inflammatory cytokines were increased in persons with cholangiocarcinoma (CCA). Different cytokine gene polymorphisms were found among patients with or without fibrosis and CCA. Interestingly, a subset of patients with the IL-6 -174G>C polymorphisms, the IFN- γ +874T>A heterozygous (TA) and homozygous (AA) variants, and the A allele of TNF- α -308G/A associated with CCA. Further studies on these gene polymorphisms and in relation to other cytokines are warranted since they may identify genotypes and phenotypes predictive of risk for CCA.

ACKNOWLEDGMENTS

This work was supported by the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission, through the Health Cluster (SHeP-GMS), Khon Kaen University, the Thailand Research Fund (TRF) grant no. RTA 5680006, the National Institute of Allergy and Infectious Diseases (NIAID), NIH, award no. P50AI098639, and National Cancer Institute (NCI), National Institutes of Health (NIH) award no. CA164719. B. Sriipa is a TRF Senior Research Scholar. The content is

solely the responsibility of the authors and does not necessarily represent the official views of the TRE, NIAID, NCI, or the NIH or the funders.

CONFLICT OF INTEREST

We have no conflict of interest related to this study.

REFERENCES

- Sripa B, Brindley PJ, Mulvenna J, Laha T, Smout MJ, Mairiang E, Bethony JM, Loukas A. The tumorigenic liver fluke *Opisthorchis viverrini*-multiple pathways to cancer. *Trends Parasitol* 2012; 28: 395-407.
- Sripa B, Kaewkes S, Sithithaworn P, Mairiang E, Laha T, Smout M, Pairojkul C, Bhudhisawasdi V, Tesana S, Thinkamrop B, Bethony JM, Loukas A, Brindley PJ. Liver fluke induces cholangiocarcinoma. *PLoS Med* 2007; 4: e201.
- Mairiang E, Chaiyakum J, Chamadol N, Laopaiboon V, Srinakaran J, Kunitipaya J, Sriamporn S, Suwanrungruang K, Vatana-sapt V. Ultrasound screening for *Opisthorchis viverrini*-associated cholangiocarcinomas: experience in an endemic area. *Asian Pac J Cancer Prev* 2006; 7: 431-433.
- Mairiang E, Laha T, Bethony JM, Thinkhamrop B, Kaewkes S, Sithithaworn P, Tesana S, Loukas A, Brindley PJ, Sripa B. Ultrasonography assessment of hepatobiliary abnormalities in 3359 subjects with *Opisthorchis viverrini* infection in endemic areas of Thailand. *Parasitol Int* 2012; 61: 208-211.
- Bradley JR. TNF-mediated inflammatory disease. *J Pathol* 2008; 214: 149-160.
- Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002; 420: 860-867.
- Ninlawan K, O'Hara SP, Splinter PL, Yongvanit P, Kaewkes S, Surapaitoon A, LaRusso NF, Sripa B. *Opisthorchis viverrini* excretory/secretory products induce toll-like receptor 4 upregulation and production of interleukin 6 and 8 in cholangiocyte. *Parasitol Int* 2010; 59: 616-621.
- Bidwell J, Keen L, Gallagher G, Kimberly R, Huizinga T, McDermott MF, Oksenberg J, McNicholl J, Pociot F, Hardt C, D'Alfonso S. Cytokine gene polymorphism in human disease: on-line databases. *Genes Immun* 1999; 1: 3-19.
- Sripa B, Mairiang E, Thinkhamrop B, Laha T, Kaewkes S, Sithithaworn P, Tessana S, Loukas A, Brindley PJ, Bethony JM. Advanced periductal fibrosis from infection with the carcinogenic human liver fluke *Opisthorchis viverrini* correlates with elevated levels of interleukin-6. *Hepatology* 2009; 50: 1273-1281.
- Surapaitoon A, Suttiprapa S, Khuntikeo N, Pairojkul C, Sripa B. Cytokine profiles in *Opisthorchis viverrini* stimulated peripheral blood mononuclear cells from cholangiocarcinoma patients. *Parasitol Int* 2016; 66: 889-892.
- Sripa B, Kaewkes S. Localisation of parasite antigens and inflammatory responses in experimental opisthorchiasis. *Int J Parasitol* 2000; 30: 735-740.
- Sripa B, Kaewkes S. Relationship between parasite-specific antibody responses and intensity of *Opisthorchis viverrini* infection in hamsters. *Parasite Immunol* 2000; 22: 139-145.
- Aida Y, Pabst MJ. Removal of endotoxin from protein solutions by phase separation using Triton X-114. *J Immunol Methods* 1990; 132: 191-195.
- Hurley JC. Endotoxemia: methods of detection and clinical correlates. *Clin Microbiol Rev* 1995; 8: 268-292.
- Duarte I, Santos A, Sousa H, Catarino R, Pinto D, Matos A, Pereira D, Moutinho J, Canedo P, Machado JC, Medeiros R. G-308A TNF-alpha polymorphism is associated with an increased risk of invasive cervical cancer. *Biochem Biophys Res Commun* 2005; 334: 588-592.
- Gatti LL, Burbano RR, Zambaldi-Tunes M, de-Lábio RW, de Assumpção PP, de Arruda Cardoso-Smith M, Marques-Payão SL. Interleukin-6 polymorphisms, *Helicobacter pylori* infection in adult Brazilian patients with chronic gastritis and gastric adenocarcinoma. *Arch Med Res* 2007; 38: 551-555.
- He B, Pan Y, Xu Y, Nie Z, Chen L, Gu L, Wang S. Increased risk for gastric cancer in carriers of the lymphotoxin-alpha+252G variant infected by *Helicobacter pylori*. *Genet Test Mol Biomarkers* 2012; 16: 9-14.
- Rad R, Prinz C, Neu B, Neuhofer M, Zeitner M, Volland P, Becker I, Schepp W, Gerhard M. Synergistic effect of *Helicobacter pylori* virulence factors and interleukin-1 polymorphisms for the development of severe histological changes in the gastric mucosa. *J Infect Dis* 2003; 188: 272-281.
- Pravica V, Perrey C, Stevens A, Lee JH, Hutchinson IV. A single nucleotide polymorphism in the first intron of the human IFN-gamma gene: absolute correlation with a polymorphic CA microsatellite marker of high IFN-gamma production. *Hum Immunol* 2000; 61: 863-866.
- Khuntikeo N, Loilome W, Thinkhamrop B, Chamadol N, Yongvanit P. A comprehensive public health conceptual framework and strategy to effectively combat cholangiocarcinoma in Thailand. *PLoS Negl Trop Dis* 2016; 10: e0004293.
- Sungkasubun P, Siripongsakun S, Akkarachinorath K, Vidhyar-korn S, Worakitsitatorn A, Sricharunrat T, Singharuksa S, Chan-wat R, Bunchaliew C, Charoenphattharaphesat S, Molek R, Yimyaem M, Sornsamdang G, Soonklang K, Wittayasak K, Aue-warakul CU, Mahidol C. Ultrasound screening for cholangiocarcinoma could detect premalignant lesions and early-stage diseases with survival benefits: a population-based prospective study of 4,225 subjects in an endemic area. *BMC Cancer* 2016; 16: 346.
- Brindley PJ, da Costa JM, Sripa B. Why does infection with some helminths cause cancer? *Trends Cancer* 2015; 1: 174-182.
- Sripa B, Thinkhamrop B, Mairiang E, Laha T, Kaewkes S, Sithithaworn P, Periago MV, Bhudhisawasdi V, Yonglithipagon P, Mulvenna J, Brindley PJ, Loukas A, Bethony JM. Elevated plasma IL-6 associates with increased risk of advanced fibrosis and chol-

- angiocarcinoma in individuals infected by *Opisthorchis viverrini*. *PLoS Negl Trop Dis* 2012; 6: e1654.
24. Chaiyadet S, Smout M, Johnson M, Whitchurch C, Turnbull L, Kaewkes S, Sotillo J, Loukas A, Sripa B. Excretory/secretory products of the carcinogenic liver fluke are endocytosed by human cholangiocytes and drive cell proliferation and IL6 production. *Int J Parasitol* 2015; 45: 773-781.
 - 25 Smout MJ, Sotillo J, Laha T, Papatpremsiri A, Rinaldi G, Pimenta RN, Chan LY, Johnson MS, Turnbull L, Whitchurch CB, Giacomini PR, Moran CS, Golledge J, Daly N, Sripa B, Mulvenna JP, Brindley PJ, Loukas A. Carcinogenic parasite secretes growth factor that accelerates wound healing and potentially promotes neoplasia. *PLoS Pathog* 2015; 11: e1005209.
 26. Cantor MJ, Nickerson P, Bernstein CN. The role of cytokine gene polymorphisms in determining disease susceptibility and phenotype in inflammatory bowel disease. *Am J Gastroenterol* 2005; 100: 1134-1142.
 27. Dai CY, Chuang WL, Hsieh MY, Lee LP, Hou NJ, Chen SC, Lin ZY, Hsieh MY, Wang LY, Tsai JF, Chang WY, Yu ML. Polymorphism of interferon-gamma gene at position +874 and clinical characteristics of chronic hepatitis C. *Transl Res* 2006; 148: 128-133.
 28. Shastry BS. SNPs: impact on gene function and phenotype. *Methods Mol Biol* 2009; 578: 3-22.
 29. Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman MM. A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science* 2007; 315: 525-528.
 30. Cordovado SK, Hendrix M, Greene CN, Mochal S, Earley MC, Farrell PM, Kharrazi M, Hannon WH, Mueller PW. CFTR mutation analysis and haplotype associations in CF patients. *Mol Genet Metab* 2012; 105: 249-254.
 31. Joshi N, Kannan S, Kotian N, Bhat S, Kale M, Hake S. Interleukin 6 -174G>C polymorphism and cancer risk: meta-analysis reveals a site dependent differential influence in Ancestral North Indians. *Hum Immunol* 2014; 75: 901-908.
 32. Wang B, Wang J, Zheng Y, Zhou S, Zheng J, Wang F, Ma X, Zeng Z. A study of TNF- α -238 and -308 polymorphisms with different outcomes of persistent hepatitis B virus infection in China. *Pathology* 2010; 42: 674-680.
 33. Kursunel MA, Esendagli G. The untold story of IFN-gamma in cancer biology. *Cytokine Growth Factor Rev* 2016; 31: 73-81.
 34. Pestka S, Krause CD, Walter MR. Interferons, interferon-like cytokines, and their receptors. *Immunol Rev* 2004; 202: 8-32.
 35. Cardoso CC, Pereira AC, Brito-de-Souza VN, Dias-Baptista IM, Maniero VC, Venturini J, Vilani-Moreno FR, de Souza FC, Ribeiro-Alves M, Sarno EN, Pacheco AG, Moraes MO. IFNG +874 T>A single nucleotide polymorphism is associated with leprosy among Brazilians. *Hum Genet* 2010; 128: 481-490.
 36. Sun Y, Lu Y, Pen Q, Li T, Xie L, Deng Y, Qin A. Interferon gamma +874 T/A polymorphism increases the risk of cervical cancer: evidence from a meta-analysis. *Tumour Biol* 2015; 36: 4555-4564.
 37. von Linsingen R, Bompeixe EP, Maestri CA, Carvalho NS, da Graca Bicalho M. IFNG (+874 T/A) polymorphism and cervical intraepithelial neoplasia in Brazilian women. *J Interferon Cytokine Res* 2009; 29: 285-288.